antibodies -online.com





MRPL53 Protein (Myc-DYKDDDDK Tag)





Go to Product page

| _ | | | |
|--------|-------|-------|-------|
| | N/0 | r\/I | ew |
| \sim | ' v C | 1 V I | C V V |

| Overview | | |
|-------------------------------|---|--|
| Quantity: | 20 μg | |
| Target: | MRPL53 | |
| Origin: | Human | |
| Source: | HEK-293 Cells | |
| Protein Type: | Recombinant | |
| Purification tag / Conjugate: | This MRPL53 protein is labelled with Myc-DYKDDDDK Tag. | |
| Application: | Antibody Production (AbP), Standard (STD) | |
| Product Details | | |
| Characteristics: | Recombinant human Mitochondrial ribosomal protein L53 (MRPL53), nuclear gene encoding mitochondrial protein protein expressed in HEK293 cells. Produced with end-sequenced ORF clone | |
| Purity: | > 80 % as determined by SDS-PAGE and Coomassie blue staining | |
| Target Details | | |
| Target: | MRPL53 | |
| Abstract: | MRPL53 Products | |
| Background: | Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75 % protein to rRNA | |

composition compared to prokaryotic ribosomes, where this ratio is reversed. Another

Target Details

difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology. This gene encodes a 39S subunit protein. A pseudogene corresponding to this gene is found on chromosome 1p.

Molecular Weight: 11.9 kDa

NCBI Accession: NP_444278

Application Details


| Application Notes: | Recombinant human proteins can be used for: | |
|--------------------|--|--|
| | Native antigens for optimized antibody production | |
| | Positive controls in ELISA and other antibody assays | |
| Comment: | The tag is located at the C-terminal. | |
| | | |

For Research Use only

Handling

Restrictions:

| Concentration: | 50 μg/mL | |
|------------------|---|--|
| Buffer: | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol. | |
| Storage: | -80 °C | |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze | |
| | immediately. Only 2-3 freeze thaw cycles are recommended. | |



Western Blotting

Image 1. Validation with Western Blot