antibodies -online.com





MNAT1 Protein (Myc-DYKDDDDK Tag)






Overview

| Overview | |
|-------------------------------|--|
| Quantity: | 20 μg |
| Target: | MNAT1 |
| Origin: | Human |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This MNAT1 protein is labelled with Myc-DYKDDDDK Tag. |
| Application: | Antibody Production (AbP), Standard (STD) |
| Product Details | |
| Characteristics: | Recombinant human MNAT1 / RNF66 protein expressed in HEK293 cells. Produced with end-sequenced ORF clone |
| Purity: | > 80 % as determined by SDS-PAGE and Coomassie blue staining |
| Target Details | |
| Target: | MNAT1 |
| Alternative Name: | Mnat1,rnf66 (MNAT1 Products) |
| Background: | The protein encoded by this gene, along with cyclin H and CDK7, forms the CDK-activating kinase (CAK) enzymatic complex. This complex activates several cyclin-associated kinases and can also associate with TFIIH to activate transcription by RNA polymerase II. Two transcript variants encoding different isoforms have been found for this gene. |
| Molecular Weight: | 35.6 kDa |

Target Details

| rarget Details | |
|---------------------|---|
| NCBI Accession: | NP_002422 |
| Pathways: | Cell Division Cycle, Mitotic G1-G1/S Phases, M Phase |
| Application Details | |
| Application Notes: | Recombinant human proteins can be used for: |
| | Native antigens for optimized antibody production |
| | Positive controls in ELISA and other antibody assays |
| Comment: | The tag is located at the C-terminal. |
| Restrictions: | For Research Use only |
| Handling | |
| Concentration: | 50 μg/mL |
| Buffer: | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol. |
| Storage: | -80 °C |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. |

Images



Western Blotting

Image 1. Validation with Western Blot