antibodies -online.com





MRPL10 Protein (His tag)





Go to Product page

| _ | | | |
|--------|-----------------------|---------|-------|
| | $ \backslash / \cap$ | r\/I | ew |
| \sim | ' V C | . I V I | C V V |

| Uverview | | |
|-------------------------------|---|--|
| Quantity: | 50 μg | |
| Target: | MRPL10 | |
| Origin: | Human | |
| Source: | Escherichia coli (E. coli) | |
| Protein Type: | Recombinant | |
| Purification tag / Conjugate: | This MRPL10 protein is labelled with His tag. | |
| Application: | Antibody Production (AbP), Standard (STD) | |
| Product Details | | |
| Characteristics: | Recombinant human MRPL10 (full length, N-term HIS tag) protein expressed in E. coli. Produced with end-sequenced ORF clone | |
| Purity: | > 80 % as determined by SDS-PAGE and Coomassie blue staining | |
| Target Details | | |
| Target: | MRPL10 | |
| Alternative Name: | Name: Mrpl10 (MRPL10 Products) | |
| Background: | Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein | |
| | synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a | |
| | small 28S subunit and a large 39S subunit. They have an estimated 75 % protein to rRNA | |
| | composition compared to prokaryotic ribosomes, where this ratio is reversed. Another | |
| | difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter | |

Target Details

| contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ |
|---|
| greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition |
| by sequence homology. This gene encodes a 39S subunit protein. Sequence analysis identified |
| three transcript variants that encode two different isoforms. A pseudogene corresponding to |
| this gene is found on chromosome 5q. |
| |

Molecular Weight:

26.3 kDa

NCBI Accession:

NP_660298

For Research Use only


Application Details

| Application Notes: | Recombinant human proteins can be used for: | |
|--------------------|--|--|
| | Native antigens for optimized antibody production | |
| | Positive controls in ELISA and other antibody assays | |
| Comment: | The tag is located at the N-terminal. | |

Handling

Restrictions:

| Concentration: | 50 μg/mL | |
|------------------|---|--|
| Buffer: | 25 mM Tris, pH 8.0, 150 mM NaCl, 10 % glycerol, 1 % Sarkosyl. Store at -80C. Avoid repeated freeze-thaw cycles. Stable for at least 3 months from receipt of products under proper storage and handling conditions. | |
| Storage: | -80 °C | |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. | |



Western Blotting

Image 1. Validation with Western Blot