
## antibodies - online.com





## MTA1 Protein (Myc-DYKDDDDK Tag)



Image



$\sim$			
Ove	rvi	ev	V

Overview	
Quantity:	20 μg
Target:	MTA1
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This MTA1 protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	<ul> <li>Recombinant human MTA1 protein expressed in HEK293 cells.</li> <li>Produced with end-sequenced ORF clone</li> </ul>
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	MTA1
Alternative Name:	Mta1 (MTA1 Products)
Background:	This gene encodes a protein that was identified in a screen for genes expressed in metastatic
	cells, specifically, mammary adenocarcinoma cell lines. Expression of this gene has been
	correlated with the metastatic potential of at least two types of carcinomas although it is also
	expressed in many normal tissues. The role it plays in metastasis is unclear. It was initially
	thought to be the 70kD component of a nucleosome remodeling deacetylase complex, NuRD,

Storage Comment:

raiget betails	
	but it is more likely that this component is a different but very similar protein. These two
	proteins are so closely related, though, that they share the same types of domains. These domains include two DNA binding domains, a dimerization domain, and a domain commonly
	found in proteins that methylate DNA. The profile and activity of this gene product suggest that
	it is involved in regulating transcription and that this may be accomplished by chromatin
	remodeling. Two transcript variants encoding different isoforms have been found for this gene.
Molecular Weight:	80.6 kDa
NCBI Accession:	NP_004680
Pathways:	Chromatin Binding
Application Details	
Application Notes:	Recombinant human proteins can be used for:
	Native antigens for optimized antibody production
	Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the C-terminal.
Restrictions:	For Research Use only
Handling	
Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C

immediately. Only 2-3 freeze thaw cycles are recommended.

Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze



## **Western Blotting**

Image 1. Validation with Western Blot