antibodies -online.com





MBP Protein (Transcript Variant 7) (Myc-DYKDDDDK Tag)



Image



Go to Product page

\sim	
()\/白	view
	V I C V V

Overview	
Quantity:	20 μg
Target:	MBP
Protein Characteristics:	Transcript Variant 7
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This MBP protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	Recombinant human Myelin Basic Protein (transcript variant 7) protein expressed in HEK293
	cells. • Produced with end-sequenced ORF clone
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	MBP
Alternative Name:	Myelin Basic Protein (MBP Products)
Background:	The protein encoded by the classic MBP gene is a major constituent of the myelin sheath of
	oligodendrocytes and Schwann cells in the nervous system. However, MBP-related transcripts
	are also present in the bone marrow and the immune system. These mRNAs arise from the

long MBP gene (otherwise called 'Golli-MBP') that contains 3 additional exons located upstream of the classic MBP exons. Alternative splicing from the Golli and the MBP transcription start sites gives rise to 2 sets of MBP-related transcripts and gene products. The Golli mRNAs contain 3 exons unique to Golli-MBP, spliced in-frame to 1 or more MBP exons. They encode hybrid proteins that have N-terminal Golli aa sequence linked to MBP aa sequence. The second family of transcripts contain only MBP exons and produce the well characterized myelin basic proteins. This complex gene structure is conserved among species suggesting that the MBP transcription unit is an integral part of the Golli transcription unit and that this arrangement is important for the function and/or regulation of these genes.

Molecular Weight:

32.9 kDa

NCBI Accession:

NP_001020272

Application Details

1 nn	lication	Motoo:
AUU	lication	MOLES.

Recombinant human proteins can be used for:

Native antigens for optimized antibody production

Positive controls in ELISA and other antibody assays

Comment:

The tag is located at the C-terminal.

Restrictions:

For Research Use only

Handling

C_{0}	วท	C	er	m	a	IJ	OI	Π

50 µg/mL

Buffer:

25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.


Storage:

-80 °C

Storage Comment:

Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze

immediately. Only 2-3 freeze thaw cycles are recommended.



Western Blotting

Image 1. Validation with Western Blot