Datasheet for ABIN2726735

# NAT15 Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)

Go to Product page

## 1 Image

## Overview

| | |
|---|---|
| Quantity: | 20 µg |
| Target: | NAT15 |
| Protein Characteristics: | Transcript Variant 1 |
| Origin: | Human |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This NAT15 protein is labelled with Myc-DYKDDDDK Tag. |
| Application: | Antibody Production (AbP), Standard (STD) |

## Product Details

| | |
|---|---|
| Characteristics: | • Recombinant human N-acetyltransferase 15 (transcript variant 1) protein expressed in HEK293 cells.<br>• Produced with end-sequenced ORF clone |
| Purity: | > 80 % as determined by SDS-PAGE and Coomassie blue staining |

## Target Details

| | |
|---|---|
| Target: | NAT15 |
| Alternative Name: | N-Acetyltransferase 15 (NAT15 Products) |
| Background: | This gene encodes an enzyme that localizes to the Golgi apparatus, where it transfers an acetyl group to the N-terminus of free proteins. This enzyme acts on histones, and its activity is important for chromatin assembly and chromosome integrity. Alternative splicing and the use |

## Target Details

| | |
|---|---|
| | of alternative promoters results in multiple transcript variants. The upstream promoter is located in a differentially methylated region (DMR) and undergoes imprinting transcript variants originating from this position are expressed from the maternal allele. |
| Molecular Weight: | 27.3 kDa |
| NCBI Accession: | NP_001077070 |

## Application Details

| | |
|---|---|
| Application Notes: | Recombinant human proteins can be used for:<br>Native antigens for optimized antibody production<br>Positive controls in ELISA and other antibody assays |
| Comment: | The tag is located at the C-terminal. |
| Restrictions: | For Research Use only |

## Handling

| | |
|---|---|
| Concentration: | 50 µg/mL |
| Buffer: | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol. |
| Storage: | -80 °C |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. |

## Images




**Western Blotting**

**Image 1.** Validation with Western Blot