antibodies -online.com





NACA Protein (Transcript Variant 4) (Myc-DYKDDDDK Tag)



Image




Go ·	to l	Prod	luct	page

Overview	
Quantity:	20 μg
Target:	NACA (NACa1)
Protein Characteristics:	Transcript Variant 4
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This NACA protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	 Recombinant human NACA (transcript variant 4) protein expressed in HEK293 cells. Produced with end-sequenced ORF clone
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	NACA (NACa1)
Alternative Name:	Naca (NACa1 Products)
Background:	This gene encodes a protein that associates with basic transcription factor 3 (BTF3) to form the nascent polypeptide-associated complex (NAC). This complex binds to nascent proteins

that lack a signal peptide motif as they emerge from the ribosome, blocking interaction with the signal recognition particle (SRP) and preventing mistranslocation to the endoplasmic reticulum.

Target Details

- al got 2 otalio		
	This protein is an IgE autoantigen in atopic dermatitis patients. Alternative splicing results in multiple transcript variants, but the full length nature of some of these variants, including those encoding very large proteins, has not been determined. There are multiple pseudogenes of this gene on different chromosomes.	
Molecular Weight:	23.2 kDa	
NCBI Accession:	NP_001106673	
Pathways:	Regulation of Muscle Cell Differentiation, Skeletal Muscle Fiber Development	
Application Details		
Application Notes:	Recombinant human proteins can be used for:	
	Native antigens for optimized antibody production	
	Positive controls in ELISA and other antibody assays	
Comment:	The tag is located at the C-terminal.	
Restrictions:	For Research Use only	
Handling		
Concentration:	50 μg/mL	
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.	
Storage:	-80 °C	
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.	



Western Blotting

Image 1. Validation with Western Blot