antibodies -online.com





NLRP2 Protein (Myc-DYKDDDDK Tag)



Image



Go to	Prod	luct	pade

U	V	eı	V	1	е	V	١

Quantity:	20 μg
Target:	NLRP2
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This NLRP2 protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	 Recombinant human NALP2 protein expressed in HEK293 cells. Produced with end-sequenced ORF clone
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	NLRP2
Alternative Name:	Nalp2 (NLRP2 Products)
Background:	This gene is a member of the nucleotide-binding and leucine-rich repeat receptor (NLR) family, and is predicted to contain an N-terminal pyrin effector domain (PYD), a centrally-located nucleotide-binding and oligomerization domain (NACHT) and C-terminal leucine-rich repeats (LRR). Members of this gene family are thought to be important regulators of immune responses. This gene product interacts with components of the IkB kinase (IKK) complex, and


Target Details

Storage Comment:

rargerbetane	
	can regulate both caspase-1 and NF-kB (nuclear factor kappa-light-chain-enhancer of activated B cells) activity. The pyrin domain is necessary and sufficient for suppression of NF-kB activity. An allelic variant (rs147585490) has been found that is incapable of blocking the transcriptional activity of NF-kB. Alternative splicing results in multiple transcript variants encoding different isoforms.
Molecular Weight:	120.3 kDa
NCBI Accession:	NP_060322
Pathways:	Production of Molecular Mediator of Immune Response, Positive Regulation of Endopeptidase Activity, Inflammasome
Application Details	
Application Notes:	Recombinant human proteins can be used for: Native antigens for optimized antibody production Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the C-terminal.
Restrictions:	For Research Use only
Handling	
Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C
0: 0 :	

immediately. Only 2-3 freeze thaw cycles are recommended.

Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze



Western Blotting

Image 1. Validation with Western Blot