antibodies.com

Datasheet for ABIN2726831 NARF Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)



1 Image

Overview

Quantity:	20 µg
Target:	NARF
Protein Characteristics:	Transcript Variant 1
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This NARF protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	 Recombinant human NARF (transcript variant 1) protein expressed in HEK293 cells. Produced with end-sequenced ORF clone
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	NARF
Alternative Name:	Narf (NARF Products)
Background:	Several proteins have been found to be prenylated and methylated at their carboxyl-terminal
	ends. Prenylation was initially believed to be important only for membrane attachment.
	However, another role for prenylation appears to be its importance in protein-protein

interactions. The only nuclear proteins known to be prenylated in mammalian cells are prelamin

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN2726831 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

	A- and B-type lamins. Prelamin A is farnesylated and carboxymethylated on the cysteine residue
	of a carboxyl-terminal CaaX motif. This post-translationally modified cysteine residue is
	removed from prelamin A when it is endoproteolytically processed into mature lamin A. The
	protein encoded by this gene binds to the prenylated prelamin A carboxyl-terminal tail domain.
	It may be a component of a prelamin A endoprotease complex. The encoded protein is located
	in the nucleus, where it partially colocalizes with the nuclear lamina. It shares limited sequence
	similarity with iron-only bacterial hydrogenases. Alternatively spliced transcript variants
	encoding different isoforms have been identified for this gene, including one with a novel exon
	that is generated by RNA editing.
Molecular Weight:	51 kDa
NCBI Accession:	NP_036468
Application Details	
Application Notes:	Recombinant human proteins can be used for:
Application Notes:	Recombinant human proteins can be used for: Native antigens for optimized antibody production
Application Notes:	Recombinant human proteins can be used for: Native antigens for optimized antibody production Positive controls in ELISA and other antibody assays
Application Notes: Comment:	Recombinant human proteins can be used for: Native antigens for optimized antibody production Positive controls in ELISA and other antibody assays The tag is located at the C-terminal.
Application Notes: Comment: Restrictions:	Recombinant human proteins can be used for: Native antigens for optimized antibody production Positive controls in ELISA and other antibody assays The tag is located at the C-terminal. For Research Use only
Application Notes: Comment: Restrictions:	Recombinant human proteins can be used for: Native antigens for optimized antibody production Positive controls in ELISA and other antibody assays The tag is located at the C-terminal. For Research Use only
Application Notes: Comment: Restrictions: Handling	Recombinant human proteins can be used for: Native antigens for optimized antibody production Positive controls in ELISA and other antibody assays The tag is located at the C-terminal. For Research Use only
Application Notes: Comment: Restrictions: Handling Concentration:	Recombinant human proteins can be used for: Native antigens for optimized antibody production Positive controls in ELISA and other antibody assays The tag is located at the C-terminal. For Research Use only 50 µg/mL
Application Notes: Comment: Restrictions: Handling Concentration: Buffer:	Recombinant human proteins can be used for:Native antigens for optimized antibody productionPositive controls in ELISA and other antibody assaysThe tag is located at the C-terminal.For Research Use only50 μg/mL25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Application Notes: Comment: Restrictions: Handling Concentration: Buffer: Storage:	Recombinant human proteins can be used for: Native antigens for optimized antibody production Positive controls in ELISA and other antibody assays The tag is located at the C-terminal. For Research Use only 50 µg/mL 50 µg/mL 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol. -80 °C
Application Notes: Comment: Restrictions: Handling Concentration: Buffer: Storage: Storage Comment:	Recombinant human proteins can be used for: Native antigens for optimized antibody production Positive controls in ELISA and other antibody assays The tag is located at the C-terminal. For Research Use only 50 µg/mL 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol. -80 °C Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze



Western Blotting

Image 1. Validation with Western Blot

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN2726831 | 09/11/2023 | Copyright antibodies-online. All rights reserved.