antibodies -online.com





NFASC Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)



Image




Go to Product page

Overview	
Quantity:	20 μg
Target:	NFASC
Protein Characteristics:	Transcript Variant 1
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This NFASC protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	 Recombinant human Neurofascin (transcript variant 1) protein expressed in HEK293 cells. Produced with end-sequenced ORF clone
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	NFASC
Alternative Name:	Neurofascin (NFASC Products)
Background:	This gene encodes an L1 family immunoglobulin cell adhesion molecule with multiple IGcam
	and fibronectin domains. The protein functions in neurite outgrowth, neurite fasciculation, and
	organization of the axon initial segment (AIS) and nodes of Ranvier on axons during early
	development. Both the AIS and nodes of Ranvier contain high densities of voltage-gated Na+

Target Details

	(Nav) channels which are clustered by interactions with cytoskeletal and scaffolding proteins including this protein, gliomedin, ankyrin 3 (ankyrin-G), and betaIV spectrin. This protein links the AIS extracellular matrix to the intracellular cytoskeleton. This gene undergoes extensive alternative splicing, and the full-length nature of some variants has not been determined.[provided by RefSeq, May 2009].
Molecular Weight:	135 kDa
NCBI Accession:	NP_001005388
Pathways:	Cell-Cell Junction Organization
Application Details	
Application Notes:	Recombinant human proteins can be used for: Native antigens for optimized antibody production Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the C-terminal.
Restrictions:	For Research Use only
Handling	
Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.



Western Blotting

Image 1. Validation with Western Blot