# antibodies -online.com







## Nkx2-2 Protein (His tag)






| Overview |
|----------|
|----------|

| Quantity:                     | 50 μg  |
|-------------------------------|--|
| Target:                       | Nkx2-2   |
| Origin:                       | Human  |
| Source:                       | Escherichia coli (E. coli)   |
| Protein Type:                 | Recombinant  |
| Purification tag / Conjugate: | This Nkx2-2 protein is labelled with His tag.  |
| Application:                  | Antibody Production (AbP), Standard (STD)  |
| Product Details               |  |
| Characteristics:              | <ul> <li>Recombinant human NKX2-2 (full length, N-term HIS tag) protein expressed in E. coli.</li> <li>Produced with end-sequenced ORF clone</li> </ul>  |
| Purity:                       | > 80 % as determined by SDS-PAGE and Coomassie blue staining   |
| Target Details                |  |
| Target:                       | Nkx2-2   |
| Alternative Name:             | Nkx2-2 (Nkx2-2 Products)   |
| Background:                   | The protein encoded by this gene contains a homeobox domain and may be involved in the morphogenesis of the central nervous system. This gene is found on chromosome 20 near NKX2-4, and these two genes appear to be duplicated on chromosome 14 in the form of TITF1 and NKX2-8. The encoded protein is likely to be a nuclear transcription factor. |
| Molecular Weight:             | 30 kDa   |
|                               |  |

#### **Target Details**

| Target Details      |  |
|---------------------|--|
| NCBI Accession:     | NP_002500  |
| Pathways:           | Dopaminergic Neurogenesis  |
| Application Details |  |
| Application Notes:  | Recombinant human proteins can be used for:  |
|                     | Native antigens for optimized antibody production  |
|                     | Positive controls in ELISA and other antibody assays   |
| Comment:            | The tag is located at the N-terminal.  |
| Restrictions:       | For Research Use only  |
| Handling            |  |
| Concentration:      | 50 μg/mL   |
| Buffer:             | 25 mM Tris, pH 8.0, 150 mM NaCl, 10 % glycerol, 1 % Sarkosyl. Store at -80C. Avoid repeated    |
|                     | freeze-thaw cycles. Stable for at least 3 months from receipt of products under proper storage |
|                     | and handling conditions.   |
| Storage:            | -80 °C   |
| Storage Comment:    | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze        |
|                     | immediately. Only 2-3 freeze thaw cycles are recommended.                                      |

#### Images



### **Western Blotting**

Image 1. Validation with Western Blot