

Datasheet for ABIN2727492

NOXA1 Protein (Myc-DYKDDDDK Tag)[Go to Product page](#)**1** Image

Overview

| | |
|-------------------------------|---|
| Quantity: | 20 µg |
| Target: | NOXA1 |
| Origin: | Human |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This NOXA1 protein is labelled with Myc-DYKDDDDK Tag. |
| Application: | Antibody Production (AbP), Standard (STD) |

Product Details

| | |
|------------------|--|
| Characteristics: | <ul style="list-style-type: none">• Recombinant human NOXA1 protein expressed in HEK293 cells.• Produced with end-sequenced ORF clone |
| Purity: | > 80 % as determined by SDS-PAGE and Coomassie blue staining |

Target Details

| | |
|-------------------|--|
| Target: | NOXA1 |
| Alternative Name: | Noxa1 (NOXA1 Products) |
| Background: | This gene encodes a protein which activates NADPH oxidases, enzymes which catalyze a reaction generating reactive oxygen species. The encoded protein contains four N-terminal tetratricopeptide domains and a C-terminal Src homology 3 domain. Interaction between the encoded protein and proteins in the oxidase regulatory complex occur via the tetratricopeptide domains. Multiple transcript variants encoding different isoforms have been found for this |

Target Details

| | |
|-------------------|---------------------------|
| | gene. |
| Molecular Weight: | 51.5 kDa |
| NCBI Accession: | NP_006638 |


Application Details

| | |
|--------------------|--|
| Application Notes: | Recombinant human proteins can be used for: Native antigens for optimized antibody production Positive controls in ELISA and other antibody assays |
| Comment: | The tag is located at the C-terminal. |
| Restrictions: | For Research Use only |

Handling

| | |
|------------------|---|
| Concentration: | 50 µg/mL |
| Buffer: | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol. |
| Storage: | -80 °C |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. |

Images



Western Blotting

Image 1. Validation with Western Blot