
# antibodies -online.com





# NR1D2 Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)



Image



Go to Product page

| _ |   |   |    |    |          |   |
|---|---|---|----|----|----------|---|
| 0 | V | 0 | r٧ | /[ | $\Theta$ | M |
|   |   |   |    |    |          |   |

| Overview                      |   |
|-------------------------------|---|
| Quantity:                     | 20 μg   |
| Target:                       | NR1D2   |
| Protein Characteristics:      | Transcript Variant 1  |
| Origin:                       | Human   |
| Source:                       | HEK-293 Cells   |
| Protein Type:                 | Recombinant   |
| Purification tag / Conjugate: | This NR1D2 protein is labelled with Myc-DYKDDDDK Tag.   |
| Application:                  | Antibody Production (AbP), Standard (STD)   |
| Product Details               |   |
| Characteristics:              | <ul> <li>Recombinant human NR1D2 (transcript variant 1) protein expressed in HEK293 cells.</li> <li>Produced with end-sequenced ORF clone</li> </ul>  |
| Purity:                       | > 80 % as determined by SDS-PAGE and Coomassie blue staining  |
| Target Details                |   |
| Target:                       | NR1D2   |
| Alternative Name:             | Nr1d2 (NR1D2 Products)  |
| Background:                   | This gene encodes a member of the nuclear hormone receptor family, specifically the NR1 subfamily of receptors. The encoded protein functions as a transcriptional repressor and may play a role in circadian rhythms and carbohydrate and lipid metabolism. Alternatively spliced transcript variants have been described. |

#### **Target Details**

| Molecular Weight:  | 64.5 kDa  |
|--|-----------|
| NCBI Accession:  | NP_005117 |
| Pathways: Nuclear Receptor Transcription Pathway, Steroid Hormone Mediated Signaling Pathway |           |


## **Application Details**

| Application Notes: | Recombinant human proteins can be used for:          |  |
|--------------------|--|--|
|                    | Native antigens for optimized antibody production    |  |
|                    | Positive controls in ELISA and other antibody assays |  |
| Comment:           | The tag is located at the C-terminal.                |  |
| Restrictions:      | For Research Use only                                |  |

### Handling

| Concentration:   | 50 μg/mL  |
|------------------|---|
| Buffer:          | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.  |
| Storage:         | -80 °C  |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. |

#### **Images**



#### **Western Blotting**

Image 1. Validation with Western Blot