antibodies -online.com





NR1H3 Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)





Go to Product page

()	11/	IN	/ie	A .
	/ // 	۱ ات	/ (−	' \/\/


Overview		
Quantity:	20 μg	
Target:	NR1H3	
Protein Characteristics:	Transcript Variant 1	
Origin:	Human	
Source:	HEK-293 Cells	
Protein Type:	Recombinant	
Purification tag / Conjugate:	This NR1H3 protein is labelled with Myc-DYKDDDDK Tag.	
Application:	Antibody Production (AbP), Standard (STD)	
Product Details		
Characteristics:	 Recombinant human NR1H3 (transcript variant 1) protein expressed in HEK293 cells. Produced with end-sequenced ORF clone 	
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining	
Target Details		
Target:	NR1H3	
Alternative Name:	Nr1h3 (NR1H3 Products)	
Background:	The protein encoded by this gene belongs to the NR1 subfamily of the nuclear receptor	
	superfamily. The NR1 family members are key regulators of macrophage function, controlling	
	transcriptional programs involved in lipid homeostasis and inflammation. This protein is highly	
	expressed in visceral organs, including liver, kidney and intestine. It forms a heterodimer with	

Target Details

rarget Details		
	retinoid X receptor (RXR), and regulates expression of target genes containing retinoid	
	response elements. Studies in mice lacking this gene suggest that it may play an important role	
	in the regulation of cholesterol homeostasis. Alternatively spliced transcript variants encoding	
	different isoforms have been found for this gene.	
Molecular Weight:	50.2 kDa	
NCBI Accession:	NP_005684	
Pathways:	Nuclear Receptor Transcription Pathway, Steroid Hormone Mediated Signaling Pathway,	
	Nuclear Hormone Receptor Binding, Cellular Response to Molecule of Bacterial Origin, Hepatitis	
	C	
Application Details		
Application Notes:	Recombinant human proteins can be used for:	
	Native antigens for optimized antibody production	
	Positive controls in ELISA and other antibody assays	
Comment:	The tag is located at the C-terminal.	
Restrictions:	For Research Use only	
Handling		
Concentration:	50 μg/mL	
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.	
Storage:	-80 °C	
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze	
	immediately. Only 2-3 freeze thaw cycles are recommended.	
Publications		
Product cited in:	Tian, Goldstein, Brown: "Insulin induction of SREBP-1c in rodent liver requires LXRα-C/EBPβ	

America, Vol. 113, Issue 29, pp. 8182-7, (2016) (PubMed).

complex." in: Proceedings of the National Academy of Sciences of the United States of



Western Blotting

Image 1. Validation with Western Blot