# antibodies -online.com





# NT5C2 Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)



Image



Go to Product page

| $\sim$ |                              |       |                |
|--------|------------------------------|-------|----------------|
| ( )    | $  \backslash / \triangle  $ | r\/I  | $\triangle VV$ |
| $\cup$ | VC                           | 1 V I | ew             |

| Quantity:                     | 20 μg  |  |
|-------------------------------|--|--|
| Target:                       | NT5C2  |  |
| Protein Characteristics:      | Transcript Variant 1   |  |
| Origin:                       | Human  |  |
| Source:                       | HEK-293 Cells  |  |
| Protein Type:                 | Recombinant  |  |
| Purification tag / Conjugate: | This NT5C2 protein is labelled with Myc-DYKDDDDK Tag.  |  |
| Application:                  | Antibody Production (AbP), Standard (STD)  |  |
| Product Details               |  |  |
| Characteristics:              | <ul> <li>Recombinant human NT5C2 (transcript variant 1) protein expressed in HEK293 cells.</li> <li>Produced with end-sequenced ORF clone</li> </ul>   |  |
| Purity:                       | > 80 % as determined by SDS-PAGE and Coomassie blue staining   |  |
| Target Details                |  |  |
| Target:                       | NT5C2  |  |
| Alternative Name:             | Nt5c2 (NT5C2 Products)   |  |
| Background:                   | May have a critical role in the maintenance of a constant composition of intracellular purine/pyrimidine nucleotides in cooperation with other nucleotidases. Preferentially hydrolyzes inosine 5'-monophosphate (IMP) and other purine nucleotides. [UniProtKB/Swiss-Prot Function] |  |

### **Target Details**

| Molecular Weight: | 64.8 kDa  |
|-------------------|-----------|
| NCBI Accession:   | NP_036361 |


# **Application Details**

| Application Notes: | Recombinant human proteins can be used for:          |  |
|--------------------|--|--|
|                    | Native antigens for optimized antibody production    |  |
|                    | Positive controls in ELISA and other antibody assays |  |
| Comment:           | The tag is located at the C-terminal.                |  |
| Restrictions:      | For Research Use only                                |  |

## Handling

| Concentration:   | 50 μg/mL  |  |
|------------------|---|--|
| Buffer:          | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.  |  |
| Storage:         | -80 °C  |  |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. |  |

#### **Images**



#### **Western Blotting**

Image 1. Validation with Western Blot