
antibodies -online.com





ORC4 Protein (Transcript Variant 3) (Myc-DYKDDDDK Tag)



Image



| _ | | | | | |
|---|---|----|---|----|---|
| U | V | er | V | Ie | W |

| Overview | |
|-------------------------------|---|
| Quantity: | 20 μg |
| Target: | ORC4 |
| Protein Characteristics: | Transcript Variant 3 |
| Origin: | Human |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This ORC4 protein is labelled with Myc-DYKDDDDK Tag. |
| Application: | Antibody Production (AbP), Standard (STD) |
| Product Details | |
| Characteristics: | Recombinant human ORC4 (transcript variant 3) protein expressed in HEK293 cells. Produced with end-sequenced ORF clone |
| Purity: | > 80 % as determined by SDS-PAGE and Coomassie blue staining |
| Target Details | |
| Target: | ORC4 |
| Alternative Name: | Orc4 (ORC4 Products) |
| Background: | The origin recognition complex (ORC) is a highly conserved six subunit protein complex essential for the initiation of the DNA replication in eukaryotic cells. Studies in yeast |
| | demonstrated that ORC binds specifically to origins of replication and serves as a platform for the assembly of additional initiation factors such as Cdc6 and Mcm proteins. This gene |

Target Details

| | encodes a subunit of the ORC complex. Several alternatively spliced transcript variants, some of which encode the same protein, have been reported for this gene. |
|-------------------|---|
| Molecular Weight: | 50.2 kDa |
| NCBI Accession: | NP_859526 |
| Pathways: | Mitotic G1-G1/S Phases, DNA Replication, Synthesis of DNA |


Application Details

| Application Notes: | Recombinant human proteins can be used for: |
|--------------------|--|
| | Native antigens for optimized antibody production |
| | Positive controls in ELISA and other antibody assays |
| Comment: | The tag is located at the C-terminal. |
| Restrictions: | For Research Use only |

Handling

| Concentration: | 50 μg/mL |
|------------------|---|
| Buffer: | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol. |
| Storage: | -80 °C |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. |

Images



Western Blotting

Image 1. Validation with Western Blot