antibodies -online.com





OXCT1 Protein (Myc-DYKDDDDK Tag)



Image



| () | 11/0 | r\ /1 | $\triangle 1 $ |
|-----|--------------------|------------|----------------|
| | $\lor \lor \vdash$ | $I \vee I$ | ew |
| | | | |

| Quantity: | 20 μg |
|-------------------------------|---|
| Target: | OXCT1 |
| Origin: | Human |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This OXCT1 protein is labelled with Myc-DYKDDDDK Tag. |
| Application: | Antibody Production (AbP), Standard (STD) |
| Product Details | |
| Characteristics: | Recombinant human OXCT1 protein expressed in HEK293 cells. Produced with end-sequenced ORF clone |
| Purity: | > 80 % as determined by SDS-PAGE and Coomassie blue staining |
| Target Details | |
| Target: | OXCT1 |
| Alternative Name: | Oxct1 (OXCT1 Products) |
| Background: | This gene encodes a member of the 3-oxoacid CoA-transferase gene family. The encoded protein is a homodimeric mitochondrial matrix enzyme that plays a central role in extrahepatic ketone body catabolism by catalyzing the reversible transfer of coenzyme A from succinyl-CoA to acetoacetate. Mutations in this gene are associated with succinyl CoA:3-oxoacid CoA transferase deficiency. |

Target Details

| Molecular Weight: | 52 kDa |
|-------------------|--|
| NCBI Accession: | NP_000427 |
| Pathways: | Positive Regulation of Peptide Hormone Secretion, Carbohydrate Homeostasis |


Application Details

| Application Notes: | Recombinant human proteins can be used for: |
|--------------------|--|
| | Native antigens for optimized antibody production |
| | Positive controls in ELISA and other antibody assays |
| Comment: | The tag is located at the C-terminal. |
| Restrictions: | For Research Use only |

Handling

| Concentration: | 50 μg/mL |
|------------------|---|
| Buffer: | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol. |
| Storage: | -80 °C |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. |

Images



Western Blotting

Image 1. Validation with Western Blot