# antibodies -online.com





## PCBP1 Protein (Myc-DYKDDDDK Tag)



Image



Go to Product page

$\sim$							
	1//	$\Box$	$r \setminus$	/ [	$\bigcirc$	1	٨,

Quantity:	20 μg	
Target:	PCBP1	
Origin:	Human	
Source:	HEK-293 Cells	
Protein Type:	Recombinant	
Purification tag / Conjugate:	This PCBP1 protein is labelled with Myc-DYKDDDDK Tag.	
Application:	Antibody Production (AbP), Standard (STD)	
Product Details		
Characteristics:	<ul> <li>Recombinant human PCBP1 protein expressed in HEK293 cells.</li> <li>Produced with end-sequenced ORF clone</li> </ul>	
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining	
Target Details		
Target:	PCBP1	
Alternative Name:	Pcbp1 (PCBP1 Products)	
Background:	This intronless gene is thought to have been generated by retrotransposition of a fully processed PCBP-2 mRNA. This gene and PCBP-2 have paralogues (PCBP3 and PCBP4) which are thought to have arisen as a result of duplication events of entire genes. The protein encoded by this gene appears to be multifunctional. It along with PCBP-2 and hnRNPK corresponds to the major cellular poly(rC)-binding protein. It contains three K-homologous (KH)	

#### **Target Details**

domains which may be involved in RNA binding. This encoded protein together with PCBP-2 also functions as translational coactivators of poliovirus RNA via a sequence-specific interaction with stem-loop IV of the IRES and promote poliovirus RNA replication by binding to its 5'-terminal cloverleaf structure. It has also been implicated in translational control of the 15-lipoxygenase mRNA, human Papillomavirus type 16 L2 mRNA, and hepatitis A virus RNA. The encoded protein is also suggested to play a part in formation of a sequence-specific alphaglobin mRNP complex which is associated with alpha-globin mRNA stability.

Molecular Weight:

37.3 kDa

NCBI Accession:

NP\_006187

#### **Application Details**

Application Notes:

Recombinant human proteins can be used for:

Native antigens for optimized antibody production

Positive controls in ELISA and other antibody assays

Comment:

The tag is located at the C-terminal.

Restrictions:

For Research Use only

#### Handling

50 μg/mL

Buffer:

25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.

Storage:

-80 °C

Storage Comment:

Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze

immediately. Only 2-3 freeze thaw cycles are recommended.



### **Western Blotting**

Image 1. Validation with Western Blot