antibodies -online.com





Datasheet for ABIN2728575

PDK2 Protein (Myc-DYKDDDDK Tag)



Image



Go to Product page

| \sim | | | | |
|--------|-----------|------|----|---|
| | $ V \cap$ | r\/I | 19 | ٨ |

| 20 μg |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| PDK2 |
| Human |
| HEK-293 Cells |
| Recombinant |
| This PDK2 protein is labelled with Myc-DYKDDDDK Tag. |
| Antibody Production (AbP), Standard (STD) |
| |
| Recombinant human PDK2 protein expressed in HEK293 cells. Produced with end-sequenced ORF clone |
| > 80 % as determined by SDS-PAGE and Coomassie blue staining |
| |
| PDK2 |
| Pdk2 (PDK2 Products) |
| This gene encodes a member of the pyruvate dehydrogenase kinase family. The encoded protein phosphorylates pyruvate dehydrogenase, down-regulating the activity of the mitochondrial pyruvate dehydrogenase complex. Overexpression of this gene may play a role in both cancer and diabetes. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. |
| |

Target Details

| Molecular Weight: | 46 kDa | |
|-------------------|----------------------------------------------------------------------------------------------------------------------------|--|
| NCBI Accession: | NP_002602 | |
| Pathways: | PI3K-Akt Signaling, RTK Signaling, Carbohydrate Homeostasis, Regulation of Carbohydrate Metabolic Process, Warburg Effect | |


Application Details

| Application Notes: | Recombinant human proteins can be used for: |
|--------------------|------------------------------------------------------|
| | Native antigens for optimized antibody production |
| | Positive controls in ELISA and other antibody assays |
| Comment: | The tag is located at the C-terminal. |
| Restrictions: | For Research Use only |

Handling

| Concentration: | 50 μg/mL |
|------------------|---------------------------------------------------------------------------------------------------------------------------------------------------|
| Buffer: | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol. |
| Storage: | -80 °C |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. |

Images



Western Blotting

Image 1. Validation with Western Blot