antibodies -online.com







PITX2 Protein (His tag)






$\Omega_{VARVIAW}$				
()) (OK) (IO) A	\sim			
	/ h	IOV	1/1/	71 A

Overview	
Quantity:	50 μg
Target:	PITX2
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Purification tag / Conjugate:	This PITX2 protein is labelled with His tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	Recombinant human PITX2 (full length, N-term HIS tag, transcript variant 2) protein expressed in E. coli.
	Produced with end-sequenced ORF clone
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	PITX2
Alternative Name:	Pitx2 (PITX2 Products)
Background:	This gene encodes a member of the RIEG/PITX homeobox family, which is in the bicoid class of
	homeodomain proteins. The encoded protein acts as a transcription factor and regulates
	procollagen lysyl hydroxylase gene expression. This protein plays a role in the terminal
	differentiation of somatotroph and lactotroph cell phenotypes, is involved in the development of

rarget Details	
	the eye, tooth and abdominal organs, and acts as a transcriptional regulator involved in basal and hormone-regulated activity of prolactin. Mutations in this gene are associated with Axenfeld-Rieger syndrome, iridogoniodysgenesis syndrome, and sporadic cases of Peters anomaly. A similar protein in other vertebrates is involved in the determination of left-right asymmetry during development. Alternatively spliced transcript variants encoding distinct isoforms have been described.
Molecular Weight:	35.2 kDa
NCBI Accession:	NP_700475
Pathways:	Retinoic Acid Receptor Signaling Pathway, Regulation of Muscle Cell Differentiation, Skeletal Muscle Fiber Development
Application Details	
Application Notes:	Recombinant human proteins can be used for: Native antigens for optimized antibody production Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the N-terminal.
Restrictions:	For Research Use only
Handling	
Concentration:	50 μg/mL
Buffer:	25 mM Tris, pH 8.0, 150 mM NaCl, 10 % glycerol, 1 % Sarkosyl. Store at -80C. Avoid repeated freeze-thaw cycles. Stable for at least 3 months from receipt of products under proper storage and handling conditions.
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze

immediately. Only 2-3 freeze thaw cycles are recommended.



Western Blotting

Image 1. Validation with Western Blot