antibodies -online.com





PLCE1 Protein (Myc-DYKDDDDK Tag)



Image




Go to Product page

\sim					
	1//	⊃r	V/I	Θ	Λ

Overview		
Quantity:	20 μg	
Target:	PLCE1	
Origin:	Human	
Source:	HEK-293 Cells	
Protein Type:	Recombinant	
Purification tag / Conjugate:	This PLCE1 protein is labelled with Myc-DYKDDDDK Tag.	
Application:	Antibody Production (AbP), Standard (STD)	
Product Details		
Characteristics:	 Recombinant human PLCE1 protein expressed in HEK293 cells. Produced with end-sequenced ORF clone 	
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining	
Target Details		
Target:	PLCE1	
Alternative Name:	Plce1 (PLCE1 Products)	
Background:	This gene encodes a phospholipase enzyme that catalyzes the hydrolysis of phosphatidylinositol-4,5-bisphosphate to generate two second messengers: inositol 1,4,5-triphosphate (IP3) and diacylglycerol (DAG). These second messengers subsequently regulate various processes affecting cell growth, differentiation, and gene expression. This enzyme is regulated by small monomeric GTPases of the Ras and Rho families and by heterotrimeric G	

Target Details

- Target Betane		
	proteins. In addition to its phospholipase C catalytic activity, this enzyme has an N-terminal domain with guanine nucleotide exchange (GEF) activity. Mutations in this gene cause early-onset nephrotic syndrome characterized by proteinuria, edema, and diffuse mesangial sclerosis or focal and segmental glomerulosclerosis. Alternative splicing results in multiple transcript variants encoding distinct isoforms.[provided by RefSeq, Sep 2009]	
Molecular Weight:	258.5 kDa	
NCBI Accession:	NP_057425	
Pathways:	EGFR Signaling Pathway, Regulation of G-Protein Coupled Receptor Protein Signaling	
Application Details		
Application Notes:	Recombinant human proteins can be used for: Native antigens for optimized antibody production Positive controls in ELISA and other antibody assays	
Comment:	The tag is located at the C-terminal.	
Restrictions:	For Research Use only	
Handling		
Concentration:	50 μg/mL	
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.	
Storage:	-80 °C	
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.	



Western Blotting

Image 1. Validation with Western Blot