
# antibodies -online.com





## PRKAA1 Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)



Overview

**Image** 



Go to Product page

Quantity:	20 μg
Target:	PRKAA1
Protein Characteristics:	Transcript Variant 1
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This PRKAA1 protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	

· Produced with end-sequenced ORF clone

• Recombinant human PRKAA1 (transcript variant 1) protein expressed in HEK293 cells.

## Target Details

Characteristics:


Purity:

Alternative Name: Prkaa1 (PRKAA1 Products)  Background: The protein encoded by this gene belongs to the ser/thr protein kinase family. It is the catalytic subunit of the 5'-prime-AMP-activated protein kinase (AMPK). AMPK is a cellular energy sensor conserved in all eukaryotic cells. The kinase activity of AMPK is activated by the stimuli that increase the cellular AMP/ATP ratio. AMPK regulates the activities of a number of key.	Target:	PRKAA1
subunit of the 5'-prime-AMP-activated protein kinase (AMPK). AMPK is a cellular energy sensor conserved in all eukaryotic cells. The kinase activity of AMPK is activated by the stimuli that	Alternative Name:	Prkaa1 (PRKAA1 Products)
	Background:	subunit of the 5'-prime-AMP-activated protein kinase (AMPK). AMPK is a cellular energy sensor

> 80 % as determined by SDS-PAGE and Coomassie blue staining

### **Target Details**

	metabolic enzymes through phosphorylation. It protects cells from stresses that cause ATP depletion by switching off ATP-consuming biosynthetic pathways. Alternatively spliced transcript variants encoding distinct isoforms have been observed.
Molecular Weight:	63.8 kDa
NCBI Accession:	NP_006242
Pathways:	AMPK Signaling, Carbohydrate Homeostasis, Regulation of Carbohydrate Metabolic Process, Warburg Effect
Application Details	
Application Notes:	Recombinant human proteins can be used for:
	Native antigens for optimized antibody production
	Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the C-terminal.
Restrictions:	For Research Use only
Handling	
Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.



### **Western Blotting**

Image 1. Validation with Western Blot