# antibodies - online.com







## PKC beta Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)



#### Image



$C \circ + \circ$	Dradi	int no	~
Go to	Prout	ıcı ba	Uŧ

( )	ve	K\ /		A .
	$\cup$	1 V/	Щ.	V۷

Overview	
Quantity:	20 μg
Target:	PKC beta (PRKCB)
Protein Characteristics:	Transcript Variant 1
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This PKC beta protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	<ul> <li>Recombinant human PRKCB (transcript variant 1) protein expressed in HEK293 cells.</li> <li>Produced with end-sequenced ORF clone</li> </ul>
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	PKC beta (PRKCB)
Alternative Name:	Prkcb (PRKCB Products)
Background:	Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a

class of tumor promoters. Each member of the PKC family has a specific expression profile
and is believed to play a distinct role in cells. The protein encoded by this gene is one of the
PKC family members. This protein kinase has been reported to be involved in many different
cellular functions, such as B cell activation, apoptosis induction, endothelial cell proliferation,
and intestinal sugar absorption. Studies in mice also suggest that this kinase may also regulate
neuronal functions and correlate fear-induced conflict behavior after stress. Alternatively
spliced transcript variants encoding distinct isoforms have been reported.


Molecular Weight:	76.7 kDa
NCBI Accession:	NP_997700
Pathways:	WNT Signaling, TCR Signaling, Thyroid Hormone Synthesis, Nuclear Hormone Receptor Binding, Chromatin Binding, Myometrial Relaxation and Contraction, VEGF Signaling, Unfolded Protein
	Response, BCR Signaling

### **Application Details**

Application Notes:	Recombinant human proteins can be used for:
	Native antigens for optimized antibody production
	Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the C-terminal.
Restrictions:	For Research Use only

#### Handling

Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze
	immediately. Only 2-3 freeze thaw cycles are recommended.



#### **Western Blotting**

Image 1. Validation with Western Blot