antibodies -online.com





PKD2 Protein (Transcript Variant 3) (Myc-DYKDDDDK Tag)



Image



Go to Product page

| O | |
|-------------------------------|---|
| Overview | |
| Quantity: | 20 μg |
| Target: | PKD2 |
| Protein Characteristics: | Transcript Variant 3 |
| Origin: | Human |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This PKD2 protein is labelled with Myc-DYKDDDDK Tag. |
| Application: | Antibody Production (AbP), Standard (STD) |
| Product Details | |
| Troduct Betaile | |
| Characteristics: | Recombinant human PRKD2 / PKD2 (transcript variant 3) protein expressed in HEK293 cells. Produced with end-sequenced ORF clone |
| Purity: | > 80 % as determined by SDS-PAGE and Coomassie blue staining |
| Target Details | |
| Target: | PKD2 |
| Alternative Name: | Prkd2,pkd2 (PKD2 Products) |
| Background: | The protein encoded by this gene belongs to the protein kinase D (PKD) family of |
| | serine/threonine protein kinases. This kinase can be activated by phorbol esters as well as by |
| | gastrin via the cholecystokinin B receptor (CCKBR) in gastric cancer cells. It can bind to |
| | diacylglycerol (DAG) in the trans-Golgi network (TGN) and may regulate basolateral membrane |
| | |

Target Details

| | protein exit from TGN. Alternative splicing results in multiple transcript variants encoding different isoforms. |
|-------------------|--|
| Molecular Weight: | 96.5 kDa |
| NCBI Accession: | NP_001073350 |
| Pathways: | cAMP Metabolic Process, Maintenance of Protein Location, Negative Regulation of Transporter Activity |


Application Details

| Application Notes: | Recombinant human proteins can be used for: |
|--------------------|--|
| | Native antigens for optimized antibody production |
| | Positive controls in ELISA and other antibody assays |
| Comment: | The tag is located at the C-terminal. |
| Restrictions: | For Research Use only |

Handling

| Concentration: | 50 μg/mL |
|------------------|---|
| Buffer: | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol. |
| Storage: | -80 °C |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. |

Images



Western Blotting

Image 1. Validation with Western Blot