antibodies -online.com





PSPC1 Protein (Myc-DYKDDDDK Tag)



Image



Go to Product page

\sim			
	$ \backslash / \square $	r\/I	$\triangle V$

Overview		
Quantity:	20 μg	
Target:	PSPC1	
Origin:	Human	
Source:	HEK-293 Cells	
Protein Type:	Recombinant	
Purification tag / Conjugate:	This PSPC1 protein is labelled with Myc-DYKDDDDK Tag.	
Application:	Antibody Production (AbP), Standard (STD)	
Product Details		
Characteristics:	Recombinant human PSPC1 / PSP1 (transcript variant alpha) protein expressed in HEK293	
	cells. • Produced with end-sequenced ORF clone	
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining	
Target Details		
Target:	PSPC1	
Alternative Name:	Pspc1,psp1 (PSPC1 Products)	
Background:	This gene encodes a nucleolar protein that localizes to punctate subnuclear structures that	
	occur close to splicing speckles, known as paraspeckles. These paraspeckles are composed of	
	RNA-protein structures that include a non-coding RNA, NEAT1/Men epsilon/beta, and the	
	Drosophila Behavior Human Splicing family of proteins, which include the product of this gene	

Target Details

and the P54NRB/NONO and PSF/SFPQ proteins. Paraspeckles may function in the control of gene expression via an RNA nuclear retention mechanism. The protein encoded by this gene is found in paraspeckles in transcriptionally active cells, but it localizes to unique cap structures at the nucleolar periphery when RNA polymerase II transcription is inhibited, or during telophase. Alternative splicing of this gene results in multiple transcript variants. A related pseudogene, which is also located on chromosome 13, has been identified.

Molecular Weight:

58.6 kDa

NCBI Accession:

NP_001035879

Application Details

Application Notes:

Recombinant human proteins can be used for:

Native antigens for optimized antibody production

Positive controls in ELISA and other antibody assays

Comment:

The tag is located at the C-terminal.

Restrictions:

For Research Use only


Handling

Concentration:	50 μg/mL	
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.	
Storage:	-80 °C	

Storage Comment:

Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze

immediately. Only 2-3 freeze thaw cycles are recommended.



Western Blotting

Image 1. Validation with Western Blot