

Datasheet for ABIN2730476

RAP1GDS1 Protein (Transcript Variant 2) (Myc-DYKDDDDK Tag)[Go to Product page](#)**1** Image

Overview

| | |
|-------------------------------|----------------------------------------------------------|
| Quantity: | 20 µg |
| Target: | RAP1GDS1 |
| Protein Characteristics: | Transcript Variant 2 |
| Origin: | Human |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This RAP1GDS1 protein is labelled with Myc-DYKDDDDK Tag. |
| Application: | Antibody Production (AbP), Standard (STD) |

Product Details

| | |
|------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Characteristics: | <ul style="list-style-type: none">• Recombinant human RAP1GDS1 (transcript variant 2) protein expressed in HEK293 cells.• Produced with end-sequenced ORF clone |
| Purity: | > 80 % as determined by SDS-PAGE and Coomassie blue staining |

Target Details

| | |
|-------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Target: | RAP1GDS1 |
| Alternative Name: | Rap1gds1 (RAP1GDS1 Products) |
| Background: | The smg GDP dissociation stimulator (smgGDS) protein is a stimulatory GDP/GTP exchange protein with GTPase activity (Riess et al., 1993 [PubMed 8262526]).[supplied by OMIM, Feb 2010]. |

Target Details

| | |
|-------------------|------------------------------------------------|
| Molecular Weight: | 66.1 kDa |
| NCBI Accession: | NP_066982 |
| Pathways: | SARS-CoV-2 Protein Interactome |


Application Details

| | |
|--------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------|
| Application Notes: | Recombinant human proteins can be used for: Native antigens for optimized antibody production Positive controls in ELISA and other antibody assays |
| Comment: | The tag is located at the C-terminal. |
| Restrictions: | For Research Use only |

Handling

| | |
|------------------|---------------------------------------------------------------------------------------------------------------------------------------------------|
| Concentration: | 50 µg/mL |
| Buffer: | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol. |
| Storage: | -80 °C |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. |

Images



Western Blotting

Image 1. Validation with Western Blot