# antibodies -online.com





## RASA1 Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)



Image



Go to Product page

$\sim$							
	1//	$rac{1}{2}$	`\/	L	Р	1	٨

Overview	
Quantity:	20 μg
Target:	RASA1
Protein Characteristics:	Transcript Variant 1
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This RASA1 protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	Recombinant human RASA1 / RasGAP (transcript variant 1) protein expressed in HEK293
	cells.  • Produced with end-sequenced ORF clone
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	RASA1
Alternative Name:	Rasa1,rasgap (RASA1 Products)
Background:	The protein encoded by this gene is located in the cytoplasm and is part of the GAP1 family of
	GTPase-activating proteins. The gene product stimulates the GTPase activity of normal RAS
	p21 but not its oncogenic counterpart. Acting as a suppressor of RAS function, the protein

enhances the weak intrinsic GTPase activity of RAS proteins resulting in the inactive GDP-
bound form of RAS, thereby allowing control of cellular proliferation and differentiation.
Mutations leading to changes in the binding sites of either protein are associated with basal cell
carcinomas. Mutations also have been associated with hereditary capillary malformations (CM)
with or without arteriovenous malformations (AVM) and Parkes Weber syndrome. Alternative
splicing results in two isoforms where the shorter isoform, lacking the N-terminal hydrophobic
region but retaining the same activity, appears to be abundantly expressed in placental but not
adult tissues.


Molecular Weight:	116.2 kDa	
NCBI Accession:	NP_002881	
Pathways:	Regulation of Actin Filament Polymerization, Signaling of Hepatocyte Growth Factor Receptor, VEGER1 Specific Signals	

### **Application Details**

Application Notes:	Recombinant human proteins can be used for:
	Native antigens for optimized antibody production
	Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the C-terminal.
Restrictions:	For Research Use only

#### Handling

Concentration:	50 μg/mL	
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.	
Storage:	-80 °C	
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze	
	immediately. Only 2-3 freeze thaw cycles are recommended.	



#### **Western Blotting**

Image 1. Validation with Western Blot