antibodies - online.com







RDH10 Protein (Myc-DYKDDDDK Tag)



Image



\sim	
()\/\p	rview
\circ	

with preference for NADP as a cofactor. Studies in mice suggest that this protein is esser for synthesis of embryonic retinoic acid and is required for limb, craniofacial, and organ development.		
Origin: Human Source: HEK-293 Cells Protein Type: Recombinant Purification tag / Conjugate: This RDH10 protein is labelled with Myc-DYKDDDDK Tag. Application: Antibody Production (AbP), Standard (STD) Product Details Characteristics: • Recombinant human RDH10 protein expressed in HEK293 cells. • Produced with end-sequenced ORF clone Purity: > 80 % as determined by SDS-PAGE and Coomassie blue staining Target Details Target: RDH10 Alternative Name: Rdh10 (RDH10 Products) Background: This gene encodes a retinol dehydrogenase, which converts all-trans-retinol to all-trans-re with preference for NADP as a cofactor. Studies in mice suggest that this protein is esser for synthesis of embryonic retinoic acid and is required for limb, craniofacial, and organ development.	Quantity:	20 μg
Source: HEK-293 Cells Protein Type: Recombinant Purification tag / Conjugate: This RDH10 protein is labelled with Myc-DYKDDDDK Tag. Application: Antibody Production (AbP), Standard (STD) Product Details Characteristics: • Recombinant human RDH10 protein expressed in HEK293 cells. • Produced with end-sequenced ORF clone Purity: > 80 % as determined by SDS-PAGE and Coomassie blue staining Target Details Target: RDH10 Alternative Name: Rdh10 (RDH10 Products) Background: This gene encodes a retinol dehydrogenase, which converts all-trans-retinol to all-trans-re with preference for NADP as a cofactor. Studies in mice suggest that this protein is esser for synthesis of embryonic retinoic acid and is required for limb, craniofacial, and organ development.	Target:	RDH10
Protein Type: Recombinant Purification tag / Conjugate: This RDH10 protein is labelled with Myc-DYKDDDDK Tag. Application: Antibody Production (AbP), Standard (STD) Product Details Characteristics: • Recombinant human RDH10 protein expressed in HEK293 cells. • Produced with end-sequenced ORF clone Purity: > 80 % as determined by SDS-PAGE and Coomassie blue staining Target Details Target: RDH10 Alternative Name: Rdh10 (RDH10 Products) Background: This gene encodes a retinol dehydrogenase, which converts all-trans-retinol to all-trans-rewith preference for NADP as a cofactor. Studies in mice suggest that this protein is essert for synthesis of embryonic retinoic acid and is required for limb, craniofacial, and organ development.	Origin:	Human
Purification tag / Conjugate: This RDH10 protein is labelled with Myc-DYKDDDDK Tag. Application: Antibody Production (AbP), Standard (STD) Product Details Characteristics: Recombinant human RDH10 protein expressed in HEK293 cells. Produced with end-sequenced ORF clone Purity: > 80 % as determined by SDS-PAGE and Coomassie blue staining Target Details Target: RDH10 Alternative Name: Rdh10 (RDH10 Products) Background: This gene encodes a retinol dehydrogenase, which converts all-trans-retinol to all-trans-retwith preference for NADP as a cofactor. Studies in mice suggest that this protein is essent for synthesis of embryonic retinoic acid and is required for limb, craniofacial, and organ development.	Source:	HEK-293 Cells
Application: Antibody Production (AbP), Standard (STD) Product Details Characteristics: Produced with end-sequenced ORF clone Purity: > 80 % as determined by SDS-PAGE and Coomassie blue staining Target Details Target: RDH10 Alternative Name: Rdh10 (RDH10 Products) Background: This gene encodes a retinol dehydrogenase, which converts all-trans-retinol to all-trans-re with preference for NADP as a cofactor. Studies in mice suggest that this protein is esser for synthesis of embryonic retinoic acid and is required for limb, craniofacial, and organ development.	Protein Type:	Recombinant
Product Details Characteristics: Recombinant human RDH10 protein expressed in HEK293 cells. Produced with end-sequenced ORF clone Purity: > 80 % as determined by SDS-PAGE and Coomassie blue staining Target Details Target: RDH10 Alternative Name: Rdh10 (RDH10 Products) Background: This gene encodes a retinol dehydrogenase, which converts all-trans-retinol to all-trans-rewith preference for NADP as a cofactor. Studies in mice suggest that this protein is essert for synthesis of embryonic retinoic acid and is required for limb, craniofacial, and organ development.	Purification tag / Conjugate:	This RDH10 protein is labelled with Myc-DYKDDDDK Tag.
Characteristics: Recombinant human RDH10 protein expressed in HEK293 cells. Produced with end-sequenced ORF clone Purity: > 80 % as determined by SDS-PAGE and Coomassie blue staining Target Details Target: RDH10 Alternative Name: Rdh10 (RDH10 Products) Background: This gene encodes a retinol dehydrogenase, which converts all-trans-retinol to all-trans-re with preference for NADP as a cofactor. Studies in mice suggest that this protein is essert for synthesis of embryonic retinoic acid and is required for limb, craniofacial, and organ development.	Application:	Antibody Production (AbP), Standard (STD)
Purity: > 80 % as determined by SDS-PAGE and Coomassie blue staining Target Details Target: RDH10 Alternative Name: Rdh10 (RDH10 Products) Background: This gene encodes a retinol dehydrogenase, which converts all-trans-retinol to all-trans-rewith preference for NADP as a cofactor. Studies in mice suggest that this protein is essert for synthesis of embryonic retinoic acid and is required for limb, craniofacial, and organ development.	Product Details	
Target Details Target: RDH10 Alternative Name: Rdh10 (RDH10 Products) Background: This gene encodes a retinol dehydrogenase, which converts all-trans-retinol to all-trans-re with preference for NADP as a cofactor. Studies in mice suggest that this protein is essent for synthesis of embryonic retinoic acid and is required for limb, craniofacial, and organ development.	Characteristics:	
Target: RDH10 Alternative Name: Rdh10 (RDH10 Products) Background: This gene encodes a retinol dehydrogenase, which converts all-trans-retinol to all-trans-re with preference for NADP as a cofactor. Studies in mice suggest that this protein is essert for synthesis of embryonic retinoic acid and is required for limb, craniofacial, and organ development.	Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Alternative Name: Rdh10 (RDH10 Products) Background: This gene encodes a retinol dehydrogenase, which converts all-trans-retinol to all-trans-re with preference for NADP as a cofactor. Studies in mice suggest that this protein is essert for synthesis of embryonic retinoic acid and is required for limb, craniofacial, and organ development.	Target Details	
Background: This gene encodes a retinol dehydrogenase, which converts all-trans-retinol to all-trans-re with preference for NADP as a cofactor. Studies in mice suggest that this protein is essent for synthesis of embryonic retinoic acid and is required for limb, craniofacial, and organ development.	Target:	RDH10
with preference for NADP as a cofactor. Studies in mice suggest that this protein is esser for synthesis of embryonic retinoic acid and is required for limb, craniofacial, and organ development.	Alternative Name:	Rdh10 (RDH10 Products)
Molecular Weight: 27.0 kDa	Background:	
Wolecular Weight. 37.9 KDa	Molecular Weight:	37.9 kDa

Target Details


Application Details

Application Notes:	Recombinant human proteins can be used for:	
	Native antigens for optimized antibody production	
	Positive controls in ELISA and other antibody assays	
Comment:	The tag is located at the C-terminal.	
Restrictions:	For Research Use only	

Handling

Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.

Images



Western Blotting

Image 1. Validation with Western Blot