## antibodies -online.com





## RGS4 Protein (Transcript Variant 2) (Myc-DYKDDDDK Tag)



Image




| C ~ .  | + ~   | 1500       | 110+     | page   |
|--------|-------|------------|----------|--------|
| ( ¬( ) | I() F | $T \cap T$ | 11 16 11 | 112016 |
|        |       |            |          |        |

| Overview                      |   |  |  |
|-------------------------------|---|--|--|
| Quantity:                     | 20 μg   |  |  |
| Target:                       | RGS4  |  |  |
| Protein Characteristics:      | Transcript Variant 2  |  |  |
| Origin:                       | Human   |  |  |
| Source:                       | HEK-293 Cells   |  |  |
| Protein Type:                 | Recombinant   |  |  |
| Purification tag / Conjugate: | This RGS4 protein is labelled with Myc-DYKDDDDK Tag.  |  |  |
| Application:                  | Antibody Production (AbP), Standard (STD)   |  |  |
| Product Details               |   |  |  |
| Characteristics:              | <ul> <li>Recombinant human RGS4 (transcript variant 2) protein expressed in HEK293 cells.</li> <li>Produced with end-sequenced ORF clone</li> </ul> |  |  |
| Purity:                       | > 80 % as determined by SDS-PAGE and Coomassie blue staining  |  |  |
| Target Details                |   |  |  |
| Target:                       | RGS4  |  |  |
| Alternative Name:             | Rgs4 (RGS4 Products)  |  |  |
| Background:                   | Regulator of G protein signaling (RGS) family members are regulatory molecules that act as  |  |  |
|                               | GTPase activating proteins (GAPs) for G alpha subunits of heterotrimeric G proteins. RGS  |  |  |
|                               | proteins are able to deactivate G protein subunits of the Gi alpha, Go alpha and Gq alpha   |  |  |
|                               | subtypes. They drive G proteins into their inactive GDP-bound forms. Regulator of G protein   |  |  |

## **Target Details**

|                     | signaling 4 belongs to this family. All RGS proteins share a conserved 120-amino acid sequence termed the RGS domain. Regulator of G protein signaling 4 protein is 37 % identical to RGS1 and 97 % identical to rat Rgs4. This protein negatively regulate signaling upstream or at the |  |  |
|---------------------|--|--|--|
|                     | level of the heterotrimeric G protein and is localized in the cytoplasm. Alternatively spliced transcript variants have been found for this gene.  |  |  |
| Molecular Weight:   | 23.1 kDa   |  |  |
| NCBI Accession:     | NP_005604  |  |  |
| Pathways:           | Myometrial Relaxation and Contraction, Regulation of G-Protein Coupled Receptor Protein Signaling  |  |  |
| Application Details |  |  |  |
| Application Notes:  | Recombinant human proteins can be used for:  Native antigens for optimized antibody production   |  |  |
|                     | Positive controls in ELISA and other antibody assays   |  |  |
| Comment:            | The tag is located at the C-terminal.  |  |  |
| Restrictions:       | For Research Use only  |  |  |
| Handling            |  |  |  |
| Concentration:      | 50 μg/mL   |  |  |
| Buffer:             | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.   |  |  |
| Storage:            | -80 °C   |  |  |
| Storage Comment:    | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.  |  |  |



## **Western Blotting**

Image 1. Validation with Western Blot