# antibodies -online.com





## RNH1 Protein (Transcript Variant 5) (Myc-DYKDDDDK Tag)



**Image** 

Alternative Name:

Background:



Go to Product page

Overview	
Quantity:	20 μg
Target:	RNH1
Protein Characteristics:	Transcript Variant 5
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This RNH1 protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	<ul> <li>Recombinant human Ribonuclease inhibitor (RNH1) (transcript variant 5) protein expressed in HEK293 cells.</li> </ul>
	Produced with end-sequenced ORF clone
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	RNH1

Placental ribonuclease inhibitor (PRI) is a member of a family of proteinaceous cytoplasmic RNase inhibitors that occur in many tissues and bind to both intracellular and extracellular

RNases (summarized by Lee et al., 1988 [PubMed 3219362]). In addition to control of

Ribonuclease Inhibitor (Rnh1) (RNH1 Products)

#### **Target Details**

intracellular RNases, the inhibitor may have a role in the regulation of angiogenin (MIM 105850). Ribonuclease inhibitor, of 50,000 Da, binds to ribonucleases and holds them in a latent form. Since neutral and alkaline ribonucleases probably play a critical role in the turnover of RNA in eukaryotic cells, RNH may be essential for control of mRNA turnover the interaction of eukaryotic cells with ribonuclease may be reversible in vivo.[supplied by OMIM, Jul 2010].

Molecular Weight: 49.8 kDa

NCBI Accession: NP\_976320

#### **Application Details**

Application Notes:

Recombinant human proteins can be used for:

Native antigens for optimized antibody production


Positive controls in ELISA and other antibody assays

The tag is located at the C-terminal.

Restrictions: For Research Use only

#### Handling

Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze
	immediately. Only 2-3 freeze thaw cycles are recommended.



### **Western Blotting**

Image 1. Validation with Western Blot