

#### Datasheet for ABIN2730920

## RPSA/Laminin Receptor Protein (Transcript Variant 2) (Myc-DYKDDDK Tag)



Go to Product pag

# 1 Image

Overview	
Quantity:	20 μg
Target:	RPSA/Laminin Receptor (RPSA)
Protein Characteristics:	Transcript Variant 2
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This RPSA/Laminin Receptor protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	<ul> <li>Recombinant human Ribosomal protein SA (RPSA), transcript variant 2 (transcript variant 2) protein expressed in HEK293 cells.</li> <li>Produced with end-sequenced ORF clone</li> </ul>
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	RPSA/Laminin Receptor (RPSA)
Abstract:	RPSA Products
Background:	Laminins, a family of extracellular matrix glycoproteins, are the major noncollagenous constituent of basement membranes. They have been implicated in a wide variety of biological processes including cell adhesion, differentiation, migration, signaling, neurite outgrowth and

metastasis. Many of the effects of laminin are mediated through interactions with cell surface
receptors. These receptors include members of the integrin family, as well as non-integrin
laminin-binding proteins. This gene encodes a high-affinity, non-integrin family, laminin receptor
1. This receptor has been variously called 67 kD laminin receptor, 37 kD laminin receptor
precursor (37LRP) and p40 ribosome-associated protein. The amino acid sequence of laminin
receptor 1 is highly conserved through evolution, suggesting a key biological function. It has
been observed that the level of the laminin receptor transcript is higher in colon carcinoma
tissue and lung cancer cell line than their normal counterparts. Also, there is a correlation
between the upregulation of this polypeptide in cancer cells and their invasive and metastatic
phenotype. Multiple copies of this gene exist, however, most of them are pseudogenes thought
to have arisen from retropositional events. Two alternatively spliced transcript variants
encoding the same protein have been found for this gene.


Molecular Weight:	32.7 kDa
NCBI Accession:	NP_001012321
Pathways:	Ribonucleoprotein Complex Subunit Organization, Ribosome Assembly

### **Application Details**

Application Notes:	Recombinant numan proteins can be used for:
	Native antigens for optimized antibody production
	Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the C-terminal.
Restrictions:	For Research Use only

#### Handling

Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze
	immediately. Only 2-3 freeze thaw cycles are recommended.



#### **Western Blotting**

Image 1. Validation with Western Blot