# antibodies -online.com







## **RUNX1T1 Protein (Transcript Variant 4) (Myc-DYKDDDDK Tag)**



#### Image



| $\sim$ | -   | $\Box$ |      | +     | page   |
|--------|-----|--------|------|-------|--------|
| ( ¬( ) | 1() | м      | ()() | 11.11 | 112(16 |
|        |     |        |      |       |        |

| Overview                      |  |  |
|-------------------------------|--|--|
| Quantity:                     | 20 μg  |  |
| Target:                       | RUNX1T1  |  |
| Protein Characteristics:      | Transcript Variant 4   |  |
| Origin:                       | Human  |  |
| Source:                       | HEK-293 Cells  |  |
| Protein Type:                 | Recombinant  |  |
| Purification tag / Conjugate: | This RUNX1T1 protein is labelled with Myc-DYKDDDDK Tag.  |  |
| Application:                  | Antibody Production (AbP), Standard (STD)  |  |
| Product Details               |  |  |
| Characteristics:              | <ul> <li>Recombinant human RUNX1T1 (transcript variant 4) protein expressed in HEK293 cells.</li> <li>Produced with end-sequenced ORF clone</li> </ul> |  |
| Purity:                       | > 80 % as determined by SDS-PAGE and Coomassie blue staining   |  |
| Target Details                |  |  |

| Target:           | RUNX1T1   |  |
|-------------------|---|--|
| Alternative Name: | Runx1t1 (RUNX1T1 Products)  |  |
| Background:       | This gene encodes a member of the myeloid translocation gene family which interact with DNA-bound transcription factors and recruit a range of corepressors to facilitate transcriptional repression. The t(821)(q22q22) translocation is one of the most frequent karyotypic abnormalities in acute myeloid leukemia. The translocation produces a chimeric gene made up |  |

#### **Target Details**

| of the 5&apos-region of the runt-related transcription factor 1 gene fused to the 3&apos-region | 1 |
|---|---|
| of this gene. The chimeric protein is thought to associate with the nuclear corepressor/histon  | Š |
| deacetylase complex to block hematopoietic differentiation. Alternative splicing results in     |   |
| multiple transcript variants.   |   |

| Molecular Weight: | 63 kDa |
|-------------------|--------|
|-------------------|--------|

NCBI Accession: NP\_783554


### **Application Details**

| Application Notes: | Recombinant human proteins can be used for:          |  |  |
|--------------------|--|--|--|
|                    | Native antigens for optimized antibody production    |  |  |
|                    | Positive controls in ELISA and other antibody assays |  |  |
| Comment:           | The tag is located at the C-terminal.                |  |  |
| Restrictions:      | For Research Use only                                |  |  |

#### Handling

| Concentration:   | 50 μg/mL  |  |
|------------------|---|--|
| Buffer:          | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.                                  |  |
| Storage:         | -80 °C  |  |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze |  |
|                  | immediately. Only 2-3 freeze thaw cycles are recommended.                               |  |

#### **Images**



#### **Western Blotting**

Image 1. Validation with Western Blot