antibodies -online.com





Shugoshin Protein (Myc-DYKDDDDK Tag)



Image



	a a ta i i a a a a paga

()	11/0	K\ /	iew	1
	\cup	ועוי	$\square \vee \vee$	ı

Quantity:	20 μg
Target:	Shugoshin (SGOL1)
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This Shugoshin protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	 Recombinant human SGOL1 / SGO1 (transcript variant B1) protein expressed in HEK293 cells.
	Produced with end-sequenced ORF clone
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	Shugoshin (SGOL1)
Alternative Name:	Sgol1,sgo1 (SGOL1 Products)
Background:	The protein encoded by this gene is a member of the shugoshin family of proteins. This protein
	is thought to protect centromeric cohesin from cleavage during mitotic prophase by preventing
	phosphorylation of a cohesin subunit. Reduced expression of this gene leads to the premature

loss of centromeric cohesion, mis-segregation of sister chromatids, and mitotic arrest.

Evidence suggests that this protein also protects a small subset of cohesin found along the length of the chromosome arms during mitotic prophase. An isoform lacking exon 6 has been shown to play a role in the cohesion of centrioles (PMID: 16582621 and PMID:18331714). Mutations in this gene have been associated with Chronic Atrial and Intestinal Dysrhythmia (CAID) syndrome, characterized by the co-occurrence of Sick Sinus Syndrome (SSS) and Chronic Intestinal Pseudo-obstruction (CIPO) within the first four decades of life (PMID:25282101). Fibroblast cells from CAID patients exhibited both increased cell proliferation and higher rates of senescence. Pseudogenes of this gene have been found on chromosomes 1 and 7. Alternative splicing results in multiple transcript variants.


Molecular Weight:	31.1 kDa
NCBI Accession:	NP_001012411
Pathways:	Maintenance of Protein Location

Application Details

Application Notes:	Recombinant human proteins can be used for:
	Native antigens for optimized antibody production
	Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the C-terminal.
Restrictions:	For Research Use only

Handling

Concentration:	50 μg/mL
Buffer: 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.	
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze
	immediately. Only 2-3 freeze thaw cycles are recommended.



Western Blotting

Image 1. Validation with Western Blot