## antibodies - online.com







## **SLIT2 Protein**



Image




$\sim$	
( )\/⊝	rview
0 1 0	VICVV

Quantity:	25 μg
Target:	SLIT2
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Application:	Antibody Production (AbP), Standard (STD), Functional Studies (Func), Protein Interaction (PI)
Product Details	
Specificity:	Optimal preservation of protein structure, post-translational modifications and functions.
Characteristics:	<ul> <li>Recombinant human SLIT2 protein expressed in HEK293 cells.</li> <li>Produced with end-sequenced ORF clone</li> </ul>
	Tested for bioactivity.
Purity:	> 95 % as determined by SDS-PAGE and Coomassie blue staining
Endotoxin Level:	Endotoxin level is <0.1 ng/μg of protein (<1EU/μg).
Biological Activity Comment:	Determined by its ability to inhibit MC3T3/E1 osteoblasts cell differentiation.
Target Details	
Target:	SLIT2
Alternative Name:	Slit2 (SLIT2 Products)

## Target Details

Background:	This gene encodes a member of the slit family of secreted glycoproteins, which are ligands for
	the Robo family of immunoglobulin receptors. Slit proteins play highly conserved roles in axon
	guidance and neuronal migration and may also have functions during other cell migration
	processes including leukocyte migration. Members of the slit family are characterized by an N-
	terminal signal peptide, four leucine-rich repeats, nine epidermal growth factor repeats, and a C
	terminal cysteine knot. Proteolytic processing of this protein gives rise to an N-terminal
	fragment that contains the four leucine-rich repeats and five epidermal growth factor repeats
	and a C-terminal fragment that contains four epidermal growth factor repeats and the cysteine
	knot. Both full length and cleaved proteins are secreted extracellularly and can function in axon
	repulsion as well as other specific processes. Alternative splicing results in multiple transcript
	variants.
Molecular Weight:	120 kDa
NCBI Accession:	NP_004778
Pathways:	Regulation of Actin Filament Polymerization, Regulation of Cell Size, Smooth Muscle Cell
	Migration
Application Details	
Application Notes:	Recombinant human proteins can be used for:
	Native antigens for optimized antibody production
	Positive controls in ELISA and other antibody assays
	Protein-protein interaction
	In vitro biochemical assays and cell-based functional assays
Restrictions:	For Research Use only
Handling	
r rannam rg	
Buffer:	Lyophilized from a 0.2 µM filtered solution of 20 mM phosphate buffer,100 mM NaCl, pH 7.2
<u> </u>	Lyophilized from a 0.2 µM filtered solution of 20 mM phosphate buffer,100 mM NaCl, pH 7.2  Resuspend the protein in the desired concentration in proper buffer
Buffer:	
Buffer: Handling Advice:	Resuspend the protein in the desired concentration in proper buffer



## **Western Blotting**

Image 1. Validation with Western Blot