# antibodies -online.com





# SMAD2 Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)



Image



Publication



Go to Product page

_					
U	V	er	V	Ie	W

Overview	
Quantity:	20 μg
Target:	SMAD2
Protein Characteristics:	Transcript Variant 1
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This SMAD2 protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	<ul> <li>Recombinant human SMAD2 (transcript variant 1) protein expressed in HEK293 cells.</li> <li>Produced with end-sequenced ORF clone</li> </ul>
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	SMAD2
Alternative Name:	Smad2 (SMAD2 Products)
Background:	The protein encoded by this gene belongs to the SMAD, a family of proteins similar to the gene
	products of the Drosophila gene 'mothers against decapentaplegic' (Mad) and the C. elegans
	gene Sma. SMAD proteins are signal transducers and transcriptional modulators that mediate
	multiple signaling pathways. This protein mediates the signal of the transforming growth factor
	gene Sma. SMAD proteins are signal transducers and transcriptional modulators that media

(TGF)-beta, and thus regulates multiple cellular processes, such as cell proliferation, apoptosis,
and differentiation. This protein is recruited to the TGF-beta receptors through its interaction
with the SMAD anchor for receptor activation (SARA) protein. In response to TGF-beta signal,
this protein is phosphorylated by the TGF-beta receptors. The phosphorylation induces the
dissociation of this protein with SARA and the association with the family member SMAD4. The
association with SMAD4 is important for the translocation of this protein into the nucleus,
where it binds to target promoters and forms a transcription repressor complex with other
cofactors. This protein can also be phosphorylated by activin type 1 receptor kinase, and
mediates the signal from the activin. Alternatively spliced transcript variants have been
observed for this gene.

Molecular Weight:	52.1 kDa
NCBI Accession:	NP_005892
Pathways:	Cell Division Cycle, Hormone Transport, Chromatin Binding, Protein targeting to Nucleus

# **Application Details**

Application Notes:	Recombinant human proteins can be used for:
	Native antigens for optimized antibody production
	Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the C-terminal.
Restrictions:	For Research Use only

# Handling

Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.


### **Publications**

Product cited in:

Atanelishvili, Shirai, Akter, Buckner, Noguchi, Silver, Bogatkevich: "M10, a caspase cleavage product of the hepatocyte growth factor receptor, interacts with Smad2 and demonstrates antifibrotic properties in vitro and in vivo." in: **Translational research: the journal of laboratory** 

and clinical medicine, Vol. 170, pp. 99-111, (2016) (PubMed).

### **Images**



### **Western Blotting**

Image 1. Validation with Western Blot