antibodies.com

Datasheet for ABIN2732220 SMAD2 Protein (Transcript Variant 2) (Myc-DYKDDDDK Tag)



Overview

Image

Quantity:	20 µg
Target:	SMAD2
Protein Characteristics:	Transcript Variant 2
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This SMAD2 protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	 Recombinant human SMAD2 (transcript variant 2) protein expressed in HEK293 cells. Produced with end-sequenced ORF clone
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	SMAD2
Alternative Name:	Smad2 (SMAD2 Products)
Background:	The protein encoded by this gene belongs to the SMAD, a family of proteins similar to the gene products of the Drosophila gene 'mothers against decapentaplegic' (Mad) and the C. elegans gene Sma. SMAD proteins are signal transducers and transcriptional modulators that mediate


multiple signaling pathways. This protein mediates the signal of the transforming growth factor

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN2732220 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

	(TGF)-beta, and thus regulates multiple cellular processes, such as cell proliferation, apoptosis,
	and differentiation. This protein is recruited to the TGF-beta receptors through its interaction
	with the SMAD anchor for receptor activation (SARA) protein. In response to TGF-beta signal,
	this protein is phosphorylated by the TGF-beta receptors. The phosphorylation induces the
	dissociation of this protein with SARA and the association with the family member SMAD4. The
	association with SMAD4 is important for the translocation of this protein into the nucleus,
	where it binds to target promoters and forms a transcription repressor complex with other
	cofactors. This protein can also be phosphorylated by activin type 1 receptor kinase, and
	mediates the signal from the activin. Alternatively spliced transcript variants have been
	observed for this gene.
Molecular Weight:	52.1 kDa
NCBI Accession:	NP_001003652
Pathways:	Cell Division Cycle, Hormone Transport, Chromatin Binding, Protein targeting to Nucleus
Application Details	
Application Notes:	Recombinant human proteins can be used for:
	Native antigens for optimized antibody production
	Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the C-terminal.
Restrictions:	For Research Use only
Handling	
Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze

immediately. Only 2-3 freeze thaw cycles are recommended.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN2732220 | 09/11/2023 | Copyright antibodies-online. All rights reserved.



Western Blotting

Image 1. Validation with Western Blot

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN2732220 | 09/11/2023 | Copyright antibodies-online. All rights reserved.