antibodies -online.com





SPG20 Protein (Transcript Variant 3) (Myc-DYKDDDDK Tag)



Image

Target Details

Alternative Name:

Background:

Target:

SPG20

Spartin,spg20 (SPG20 Products)



Go to Product page

Overview	
Quantity:	20 μg
Target:	SPG20
Protein Characteristics:	Transcript Variant 3
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This SPG20 protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	 Recombinant human Spartin / SPG20 (transcript variant 3) protein expressed in HEK293 cells. Produced with end-sequenced ORF clone
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
T D !!	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn | International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com | Page 1/3 | Product datasheet for ABIN2732509 | 01/16/2024 | Copyright antibodies-online. All rights reserved.


This gene encodes a protein containing a MIT (Microtubule Interacting and Trafficking

molecule) domain, and is implicated in regulating endosomal trafficking and mitochondria

function. The protein localizes to mitochondria and partially co-localizes with microtubules.

Target Details

Target Details	
	Stimulation with epidermal growth factor (EGF) results in protein translocation to the plasma membrane, and the protein functions in the degradation and intracellular trafficking of EGF receptor. Multiple alternatively spliced variants, encoding the same protein, have been identified. Mutations associated with this gene cause autosomal recessive spastic paraplegia 20 (Troyer syndrome).
Molecular Weight:	72.7 kDa
NCBI Accession:	NP_001135767
Pathways:	Regulation of Cell Size, SARS-CoV-2 Protein Interactome
Application Details	
Application Notes:	Recombinant human proteins can be used for: Native antigens for optimized antibody production Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the C-terminal.
Restrictions:	For Research Use only
Handling	
Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.



Western Blotting

Image 1. Validation with Western Blot