antibodies -online.com





SPHK1 Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)



Image




| Go to | D | 1 | |
|-------|------|----------|------|
| | Pron | וויאו וו | nane |
| | | | |

| Overview | |
|-------------------------------|---|
| Quantity: | 20 μg |
| Target: | SPHK1 |
| Protein Characteristics: | Transcript Variant 1 |
| Origin: | Human |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This SPHK1 protein is labelled with Myc-DYKDDDDK Tag. |
| Application: | Antibody Production (AbP), Standard (STD) |
| Product Details | |
| Characteristics: | Recombinant human Sphingosine kinase 1 (SPHK1) (transcript variant 1) protein expressed in HEK293 cells. Produced with end-sequenced ORF clone |
| Purity: | > 80 % as determined by SDS-PAGE and Coomassie blue staining |
| Target Details | |
| Target: | SPHK1 |
| Abstract: | SPHK1 Products |
| Background: | The protein encoded by this gene catalyzes the phosphorylation of sphingosine to form sphingosine-1-phosphate (S1P), a lipid mediator with both intra- and extracellular functions. |

Intracellularly, S1P regulates proliferation and survival, and extracellularly, it is a ligand for cell

Target Details

| g | | |
|---------------------|---|--|
| | surface G protein-coupled receptors. This protein, and its product S1P, play a key role in TNF-alpha signaling and the NF-kappa-B activation pathway important in inflammatory, antiapoptotic, and immune processes. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. | |
| Molecular Weight: | 43.8 kDa | |
| NCBI Accession: | NP_068807 | |
| Pathways: | VEGF Signaling | |
| Application Details | | |
| Application Notes: | Recombinant human proteins can be used for: Native antigens for optimized antibody production Positive controls in ELISA and other antibody assays | |
| Comment: | The tag is located at the C-terminal. | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Concentration: | 50 μg/mL | |
| Buffer: | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol. | |
| Storage: | -80 °C | |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. | |



Western Blotting

Image 1. Validation with Western Blot