

Datasheet for ABIN2732767

**STARD5 Protein (Myc-DYKDDDDK Tag)**[Go to Product page](#)**1** Image

## Overview

Quantity:	20 µg
Target:	STARD5
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This STARD5 protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)

## Product Details

Characteristics:	<ul style="list-style-type: none"><li>• Recombinant human STARD5 protein expressed in HEK293 cells.</li><li>• Produced with end-sequenced ORF clone</li></ul>
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining

## Target Details

Target:	STARD5
Alternative Name:	Stard5 ( <a href="#">STARD5 Products</a> )
Background:	Proteins containing a steroidogenic acute regulatory-related lipid transfer (START) domain are often involved in the trafficking of lipids and cholesterol between diverse intracellular membranes. This gene is a member of the StarD subfamily that encodes START-related lipid transfer proteins. The protein encoded by this gene is a cholesterol transporter and is also able to bind and transport other sterol-derived molecules related to the cholesterol/bile acid

## Target Details

biosynthetic pathways such as 25-hydroxycholesterol. Its expression is upregulated during endoplasmic reticulum (ER) stress. The protein is thought to act as a cytosolic sterol transporter that moves cholesterol between intracellular membranes such as from the cytoplasm to the ER and from the ER to the Golgi apparatus. Alternative splicing of this gene produces multiple transcript variants.

Molecular Weight: 23.6 kDa

NCBI Accession: [NP\\_871629](#)

Pathways: [Metabolism of Steroid Hormones and Vitamin D](#), [C21-Steroid Hormone Metabolic Process](#)

## Application Details

Application Notes: Recombinant human proteins can be used for:  
Native antigens for optimized antibody production  
Positive controls in ELISA and other antibody assays

Comment: The tag is located at the C-terminal.

Restrictions: For Research Use only


## Handling

Concentration: 50 µg/mL

Buffer: 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.

Storage: -80 °C

Storage Comment: Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.



Western Blotting

**Image 1.** Validation with Western Blot