antibodies -online.com





STAT2 Protein (Myc-DYKDDDDK Tag)



Image




Go to	D		
	Pron	וויאווו	nane

Overview	
Quantity:	20 μg
Target:	STAT2
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This STAT2 protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	 Recombinant human STAT2 protein expressed in HEK293 cells. Produced with end-sequenced ORF clone
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	STAT2
Alternative Name:	Stat2 (STAT2 Products)
Background:	The protein encoded by this gene is a member of the STAT protein family. In response to
	cytokines and growth factors, STAT family members are phosphorylated by the receptor
	associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus
	where they act as transcription activators. In response to interferon (IFN), this protein forms a
	complex with STAT1 and IFN regulatory factor family protein p48 (ISGF3G), in which this

Target Details

	protein acts as a transactivator, but lacks the ability to bind DNA directly. Transcription adaptor P300/CBP (EP300/CREBBP) has been shown to interact specifically with this protein, which is thought to be involved in the process of blocking IFN-alpha response by adenovirus. Multiple transcript variants encoding different isoforms have been found for this gene.	
Molecular Weight:	97.7 kDa	
NCBI Accession:	NP_005410	
Pathways:	JAK-STAT Signaling, Hepatitis C, CXCR4-mediated Signaling Events	
Application Details		
Application Notes:	Recombinant human proteins can be used for: Native antigens for optimized antibody production Positive controls in ELISA and other antibody assays	
Comment:	The tag is located at the C-terminal.	
Restrictions:	For Research Use only	
Handling		
Concentration:	50 μg/mL	
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.	
Storage:	-80 °C	
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.	



Western Blotting

Image 1. Validation with Western Blot