antibodies - online.com





STAT6 Protein (Myc-DYKDDDDK Tag)



Image




\sim				
	$ V \cap$	r\/I	19	٨

Overview		
Quantity:	20 μg	
Target:	STAT6	
Origin:	Human	
Source:	HEK-293 Cells	
Protein Type:	Recombinant	
Purification tag / Conjugate:	This STAT6 protein is labelled with Myc-DYKDDDDK Tag.	
Application:	Antibody Production (AbP), Standard (STD)	
Product Details		
Characteristics:	 Recombinant human STAT6 protein expressed in HEK293 cells. Produced with end-sequenced ORF clone 	
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining	
Target Details		
Target:	STAT6	
Alternative Name:	Stat6 (STAT6 Products)	
Background:	The protein encoded by this gene is a member of the STAT family of transcription factors. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein plays a central role in exerting IL4 mediated biological responses. It is found to induce the expression of BCL2L1/BCL-X(L),	

Target Details

	which is responsible for the anti-apoptotic activity of IL4. Knockout studies in mice suggested the roles of this gene in differentiation of T helper 2 (Th2) cells, expression of cell surface markers, and class switch of immunoglobulins. Alternative splicing results in multiple transcript variants.[provided by RefSeq, May 2010]	
Molecular Weight:	94 kDa	
NCBI Accession:	NP_003144	
Pathways:	JAK-STAT Signaling, Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process, Production of Molecular Mediator of Immune Response	
Application Details		
Application Notes:	Recombinant human proteins can be used for: Native antigens for optimized antibody production Positive controls in ELISA and other antibody assays	
Comment:	The tag is located at the C-terminal.	
Restrictions:	For Research Use only	
Handling		
Concentration:	50 μg/mL	
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.	
Storage:	-80 °C	
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.	



Western Blotting

Image 1. Validation with Western Blot