antibodies -online.com





SULT1C2 Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)



Image



Go to Product page

| \sim | | | | | |
|--------|-----|----|----|-------------|---|
| | 1// | r۱ | /1 | \triangle | ٨ |

| Overview | |
|-------------------------------|--|
| Quantity: | 20 μg |
| Target: | SULT1C2 |
| Protein Characteristics: | Transcript Variant 1 |
| Origin: | Human |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This SULT1C2 protein is labelled with Myc-DYKDDDDK Tag. |
| Application: | Antibody Production (AbP), Standard (STD) |
| Product Details | |
| Characteristics: | Recombinant human SULT1C2 / SULT1C1 (transcript variant 1) protein expressed in HEK293 |
| | cells. • Produced with end-sequenced ORF clone |
| Purity: | > 80 % as determined by SDS-PAGE and Coomassie blue staining |
| Target Details | |
| Target: | SULT1C2 |
| Alternative Name: | Sult1c2,sult1c1 (SULT1C2 Products) |
| Background: | Sulfotransferase enzymes catalyze the sulfate conjugation of many hormones, |
| | neurotransmitters, drugs, and xenobiotic compounds. These cytosolic enzymes are different in |
| | their tissue distributions and substrate specificities. The gene structure (number and length of |

Target Details

| exons) is similar among family members. This gene encodes a protein that belongs to the |
|---|
| SULT1 subfamily, responsible for transferring a sulfo moiety from PAPS to phenol-containing |
| compounds. Two alternatively spliced transcript variants encoding different isoforms have |
| been described for this gene. |

| Molecular Weight: | 34.7 kDa |
|-------------------|----------|
|-------------------|----------|

NCBI Accession: NP_001047


Application Details

| Application Notes: | Recombinant human proteins can be used for: | |
|--------------------|--|--|
| | Native antigens for optimized antibody production | |
| | Positive controls in ELISA and other antibody assays | |
| Comment: | The tag is located at the C-terminal. | |
| Restrictions: | For Research Use only | |

Handling

| Concentration: | 50 μg/mL | |
|--|--|--|
| Buffer: | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol. | |
| Storage: | -80 °C | |
| Storage Comment: Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. | | |

Images



Western Blotting

Image 1. Validation with Western Blot