antibodies .- online.com





SUMO2 Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)



Overview

Image



Go to Product page

| Quantity: | 20 μg |
|-------------------------------|---|
| Target: | SUM02 |
| Protein Characteristics: | Transcript Variant 1 |
| Origin: | Human |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This SUMO2 protein is labelled with Myc-DYKDDDDK Tag. |
| Application: | Antibody Production (AbP), Standard (STD) |

Product Details

Characteristics:

| Purity: | > 80 % as determined by SDS-PAGE and Coomassie blue staining |
|---------------------------|--|
| Target Details | |
| | |
| Target: | SUM02 |
| Target: Alternative Name: | SUM02 Sumo2 (SUM02 Products) |

· Produced with end-sequenced ORF clone

• Recombinant human SUMO2 (transcript variant 1) protein expressed in HEK293 cells.

protein family. It functions in a manner similar to ubiquitin in that it is bound to target proteins

as part of a post-translational modification system. However, unlike ubiquitin which targets proteins for degradation, this protein is involved in a variety of cellular processes, such as

Target Details

| - al got 2 otalio | |
|---------------------|---|
| | nuclear transport, transcriptional regulation, apoptosis, and protein stability. It is not active until the last two amino acids of the carboxy-terminus have been cleaved off. Numerous pseudogenes have been reported for this gene. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. |
| Molecular Weight: | 10.7 kDa |
| NCBI Accession: | NP_008868 |
| Pathways: | Methionine Biosynthetic Process |
| Application Details | |
| Application Notes: | Recombinant human proteins can be used for: |
| | Native antigens for optimized antibody production |
| | Positive controls in ELISA and other antibody assays |
| Comment: | The tag is located at the C-terminal. |
| Restrictions: | For Research Use only |
| Handling | |
| Concentration: | 50 μg/mL |
| Buffer: | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol. |
| Storage: | -80 °C |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. |



Western Blotting

Image 1. Validation with Western Blot