
antibodies -online.com





TAPBP Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)



Overview

Purity:

Target Details

Image



Quantity:	20 μg
Target:	TAPBP
Protein Characteristics:	Transcript Variant 1
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This TAPBP protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	Recombinant human Tapasin (TAPBP) (transcript variant 1) protein expressed in HEK293 cells.
	Produced with end-sequenced ORF clone

Target: TAPBP Alternative Name: Tapasin (Tapbp) (TAPBP Products) Background: This gene encodes a transmembrane glycoprotein which mediates interaction between newly assembled major histocompatibility complex (MHC) class I molecules and the transporter associated with antigen processing (TAP), which is required for the transport of antigenic

> 80 % as determined by SDS-PAGE and Coomassie blue staining

Target Details

peptides across the endoplasmic reticulum membrane. This interaction is essential for optimal peptide loading on the MHC class I molecule. Up to four complexes of MHC class I and this protein may be bound to a single TAP molecule. This protein contains a C-terminal double-lysine motif (KKKAE) known to maintain membrane proteins in the endoplasmic reticulum. This gene lies within the major histocompatibility complex on chromosome 6. Alternative splicing results in three transcript variants encoding different isoforms.

Molecular Weight:

45.6 kDa

NCBI Accession:

NP_003181

Application Details

Application Notes:

Recombinant human proteins can be used for:

Native antigens for optimized antibody production

Positive controls in ELISA and other antibody assays

Comment:

The tag is located at the C-terminal.

Restrictions:

For Research Use only

Handling

Concentration:	

 $50 \, \mu g/mL$

Buffer:

25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.


Storage:

-80 °C

Storage Comment:

Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze

immediately. Only 2-3 freeze thaw cycles are recommended.



Western Blotting

Image 1. Validation with Western Blot