antibodies -online.com





TXNRD1 Protein (Transcript Variant 3) (Myc-DYKDDDDK Tag)



Image



Go to	Prod	uct	page

\sim							
0	۱۱/	Δ	r\	/ I		1/	١.
\cup	v	$\overline{}$	ΙV	1	$\overline{}$	٧	٧

Overview	
Quantity:	20 μg
Target:	TXNRD1
Protein Characteristics:	Transcript Variant 3
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This TXNRD1 protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	Recombinant human Thioredoxin reductase 1 / TXNRD1 (transcript variant 3) protein
	expressed in HEK293 cells. • Produced with end-sequenced ORF clone
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	TXNRD1
Alternative Name:	Thioredoxin Reductase 1,txnrd1 (TXNRD1 Products)
Background:	This gene encodes a member of the family of pyridine nucleotide oxidoreductases. This protein
	reduces thioredoxins as well as other substrates, and plays a role in selenium metabolism and
	protection against oxidative stress. The functional enzyme is thought to be a homodimer which

uses FAD as a cofactor. Each subunit contains a selenocysteine (Sec) residue which is required
for catalytic activity. The selenocysteine is encoded by the UGA codon that normally signals
translation termination. The 3' UTR of selenocysteine-containing genes have a common stem-
loop structure, the sec insertion sequence (SECIS), that is necessary for the recognition of UGA
as a Sec codon rather than as a stop signal. Alternative splicing results in several transcript
variants encoding the same or different isoforms.


Molecular Weight:	54.6 kDa
NCBI Accession:	NP_877393
Pathways:	Regulation of Lipid Metabolism by PPARalpha, Regulation of Carbohydrate Metabolic Process,
	Cell RedoxHomeostasis

Application Details

Application Notes:	Recombinant human proteins can be used for:
	Native antigens for optimized antibody production
	Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the C-terminal.
Restrictions:	For Research Use only

Handling

Concentration:	50 μg/mL	
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.	
Storage:	-80 °C	
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.	



Western Blotting

Image 1. Validation with Western Blot