antibodies -online.com





TYROBP Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)



Image

Target Details



Go to Product page

Overview	
Quantity:	20 μg
Target:	TYROBP
Protein Characteristics:	Transcript Variant 1
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This TYROBP protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	Recombinant human TYROBP / DAP12 (transcript variant 1) protein expressed in HEK293 cells.
	Produced with end-sequenced ORF clone
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining

Target:	TYROBP
Alternative Name:	Tyrobp,dap12 (TYROBP Products)
Background:	This gene encodes a transmembrane signaling polypeptide which contains an immunoreceptor tyrosine-based activation motif (ITAM) in its cytoplasmic domain. The encoded protein may associate with the killer-cell inhibitory receptor (KIR) family of membrane glycoproteins and

may act as an activating signal transduction element. This protein may bind zeta-chain (TCR) associated protein kinase 70 kDa (ZAP-70) and spleen tyrosine kinase (SYK) and play a role in signal transduction, bone modeling, brain myelination, and inflammation. Mutations within this gene have been associated with polycystic lipomembranous osteodysplasia with sclerosing leukoencephalopathy (PLOSL), also known as Nasu-Hakola disease. Its putative receptor, triggering receptor expressed on myeloid cells 2 (TREM2), also causes PLOSL. Multiple alternative transcript variants encoding distinct isoforms have been identified for this gene.

Molecular Weight:

9.3 kDa

NCBI Accession:

NP_003323

Application Details

Application Notes:

Recombinant human proteins can be used for:

Native antigens for optimized antibody production

Positive controls in ELISA and other antibody assays

Comment:

The tag is located at the C-terminal.

Restrictions:

For Research Use only

Handling

Concentration:

50 μg/mL

Buffer:

25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.


Storage:

-80 °C

Storage Comment:

Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze

immediately. Only 2-3 freeze thaw cycles are recommended.



Western Blotting

Image 1. Validation with Western Blot