## antibodies -online.com





## ube3a Protein (Transcript Variant 3) (Myc-DYKDDDDK Tag)



Image



Go to	D	1	
	Pron	וויאו וו	nane

Overview	
Quantity:	20 μg
Target:	ube3a
Protein Characteristics:	Transcript Variant 3
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This ube3a protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	<ul> <li>Recombinant human UBE3A (transcript variant 3) protein expressed in HEK293 cells.</li> <li>Produced with end-sequenced ORF clone</li> </ul>
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	ube3a
Alternative Name:	Ube3a (ube3a Products)
Background:	This gene encodes an E3 ubiquitin-protein ligase, part of the ubiquitin protein degradation system. This imprinted gene is maternally expressed in brain and biallelically expressed in other tissues. Maternally inherited deletion of this gene causes Angelman Syndrome, characterized
	by severe motor and intellectual retardation, ataxia, hypotonia, epilepsy, absence of speech, and


## **Target Details**

Storage Comment:

	characteristic facies. The protein also interacts with the E6 protein of human papillomavirus types 16 and 18, resulting in ubiquitination and proteolysis of tumor protein p53. Alternative splicing of this gene results in three transcript variants encoding three isoforms with different N-termini. Additional transcript variants have been described, but their full length nature has not been determined.
Molecular Weight:	99.9 kDa
NCBI Accession:	NP_570854
Pathways:	Intracellular Steroid Hormone Receptor Signaling Pathway
Application Details	
Application Notes:	Recombinant human proteins can be used for:
	Native antigens for optimized antibody production  Positive controls in ELISA and other antibody assays
Comment:	
Comment:  Restrictions:	Positive controls in ELISA and other antibody assays
	Positive controls in ELISA and other antibody assays  The tag is located at the C-terminal.
Restrictions:	Positive controls in ELISA and other antibody assays  The tag is located at the C-terminal.
Restrictions: Handling	Positive controls in ELISA and other antibody assays  The tag is located at the C-terminal.  For Research Use only

Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze

immediately. Only 2-3 freeze thaw cycles are recommended.



## **Western Blotting**

Image 1. Validation with Western Blot