antibodies -online.com





Ubiquilin 1 Protein (UBQLN1) (Transcript Variant 2) (Myc-DYKDDDK Tag)



Go to Product pag

1 Image

| Overview | |
|-------------------------------|---|
| Quantity: | 20 μg |
| Target: | Ubiquilin 1 (UBQLN1) |
| Protein Characteristics: | Transcript Variant 2 |
| Origin: | Human |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This Ubiquilin 1 protein is labelled with Myc-DYKDDDDK Tag. |
| Application: | Antibody Production (AbP), Standard (STD) |
| Product Details | |
| Characteristics: | Recombinant human Ubiquilin-1 (UBQLN1) (transcript variant 2) protein expressed in |
| | HEK293 cells.Produced with end-sequenced ORF clone |
| Purity: | > 80 % as determined by SDS-PAGE and Coomassie blue staining |
| Target Details | |
| Target: | Ubiquilin 1 (UBQLN1) |
| Alternative Name: | Ubiquilin-1 (Ubqln1) (UBQLN1 Products) |
| Background: | This gene encodes an ubiquitin-like protein (ubiquilin) that shares a high degree of similarity |
| | with related products in yeast, rat and frog. Ubiquilins contain an N-terminal ubiquitin-like |
| | domain and a C-terminal ubiquitin-associated domain. They physically associate with both |

Target Details

proteasomes and ubiquitin ligases, and thus are thought to functionally link the ubiquitination machinery to the proteasome to affect in vivo protein degradation. This ubiquilin has also been shown to modulate accumulation of presenilin proteins, and it is found in lesions associated with Alzheimer&aposs and Parkinson&aposs disease. Two transcript variants encoding different isoforms have been found for this gene.

Molecular Weight:

59 kDa

NCBI Accession:

NP_444295

Application Details

Application Notes: Recombinant human proteins can be used for:

Native antigens for optimized antibody production


Positive controls in ELISA and other antibody assays

Comment: The tag is located at the C-terminal.

Restrictions: For Research Use only

Handling

| Concentration: | 50 μg/mL |
|------------------|---|
| Buffer: | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol. |
| Storage: | -80 °C |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. |



Western Blotting

Image 1. Validation with Western Blot