# antibodies -online.com





# **UCK1 Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)**





| Overview                      |   |
|-------------------------------|---|
| Quantity:                     | 20 μg   |
| Target:                       | UCK1  |
| Protein Characteristics:      | Transcript Variant 1  |
| Origin:                       | Human   |
| Source:                       | HEK-293 Cells   |
| Protein Type:                 | Recombinant   |
| Purification tag / Conjugate: | This UCK1 protein is labelled with Myc-DYKDDDDK Tag.  |
| Application:                  | Antibody Production (AbP), Standard (STD)   |
| Product Details               |   |
| Characteristics:              | <ul> <li>Recombinant human UCK1 (transcript variant 1) protein expressed in HEK293 cells.</li> <li>Produced with end-sequenced ORF clone</li> </ul>   |
| Purity:                       | > 80 % as determined by SDS-PAGE and Coomassie blue staining  |
| Target Details                |   |
| Target:                       | UCK1  |
| Alternative Name:             | Uck1 (UCK1 Products)  |
| Background:                   | This gene encodes a uridine-cytidine kinase that catalyzes the phosphorylation of uridine and cytidine to uridine monophosphate (UMP) and cytidine monophosphate (CMP) but not the phosphorylation of deoxyribonucleosides or purine ribonucleosides. This enzyme can also phosphorylate uridine and cytidine analogs and uses both ATP and GTP as a phosphate donor. |

#### **Target Details**

|                   | Alternative splicing results in multiple splice variants encoding distinct isoforms. |
|-------------------|--|
| Molecular Weight: | 31.3 kDa   |
| NCBI Accession:   | NP_113620  |
| Pathways:         | Ribonucleoside Biosynthetic Process  |


## **Application Details**

| Application Notes: | Recombinant human proteins can be used for:          |
|--------------------|--|
|                    | Native antigens for optimized antibody production    |
|                    | Positive controls in ELISA and other antibody assays |
| Comment:           | The tag is located at the C-terminal.                |
| Restrictions:      | For Research Use only                                |

## Handling

| Concentration:   | 50 μg/mL  |
|------------------|---|
| Buffer:          | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.  |
| Storage:         | -80 °C  |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. |

#### **Images**



#### **Western Blotting**

Image 1. Validation with Western Blot