antibodies -online.com





UCK2 Protein (Myc-DYKDDDDK Tag)



Image



Go to Product page

\sim			
	N/6	1//r	$I \cap V$

20 μg	
UCK2	
Human	
HEK-293 Cells	
Recombinant	
This UCK2 protein is labelled with Myc-DYKDDDDK Tag.	
Antibody Production (AbP), Standard (STD)	
 Recombinant human UCK2 protein expressed in HEK293 cells. Produced with end-sequenced ORF clone 	
> 80 % as determined by SDS-PAGE and Coomassie blue staining	
UCK2	
Uck2 (UCK2 Products)	
Phosphorylates uridine and cytidine to uridine monophosphate and cytidine monophosphate. Does not phosphorylate deoxyribonucleosides or purine ribonucleosides. Can use ATP or GTP as a phosphate donor. Can also phosphorylate cytidine and uridine nucleoside analogs such as 6-azauridine, 5-fluorouridine, 4-thiouridine, 5-bromouridine, N(4)-acetylcytidine, N(4)-benzoylcytidine, 5-fluorocytidine, 2-thiocytidine, 5-methylcytidine, and N(4)-anisoylcytidine.	

Target Details

	[UniProtKB/Swiss-Prot Function]	
Molecular Weight:	29.1 kDa	
NCBI Accession:	NP_036606	
Pathways:	Ribonucleoside Biosynthetic Process, Feeding Behaviour	


Application Details

Application Notes:	Recombinant human proteins can be used for:	
	Native antigens for optimized antibody production	
	Positive controls in ELISA and other antibody assays	
Comment:	The tag is located at the C-terminal.	
Restrictions:	For Research Use only	

Handling

Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.

Images



Western Blotting

Image 1. Validation with Western Blot